

Goat Anti-Hamartin / TSC1 (isoform 1) Antibody
Peptide-affinity purified goat antibody
Catalog # AF1519a

Specification

Goat Anti-Hamartin / TSC1 (isoform 1) Antibody - Product Information

Application	WB, IHC
Primary Accession	O92574
Other Accession	NP_000359 , 7248 , 64930 (mouse)
Reactivity	Human
Predicted	Mouse, Rat, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	129767

Goat Anti-Hamartin / TSC1 (isoform 1) Antibody - Additional Information

Gene ID 7248

Other Names

Hamartin, Tuberous sclerosis 1 protein, TSC1, KIAA0243, TSC

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-Hamartin / TSC1 (isoform 1) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-Hamartin / TSC1 (isoform 1) Antibody - Protein Information

Name TSC1 {ECO:0000303|PubMed:9242607, ECO:0000312|HGNC:HGNC:12362}

Function

Non-catalytic component of the TSC-TBC complex, a multiprotein complex that acts as a negative regulator of the canonical mTORC1 complex, an evolutionarily conserved central nutrient sensor that stimulates anabolic reactions and macromolecule biosynthesis to promote cellular biomass generation and growth (PubMed:12172553, PubMed:12271141, PubMed:12906785)

target="_blank">12906785, PubMed:15340059, PubMed:24529379, PubMed:28215400). The TSC-TBC complex acts as a GTPase-activating protein (GAP) for the small GTPase RHEB, a direct activator of the protein kinase activity of mTORC1 (PubMed:12906785, PubMed:15340059, PubMed:24529379). In absence of nutrients, the TSC-TBC complex inhibits mTORC1, thereby preventing phosphorylation of ribosomal protein S6 kinase (RPS6KB1 and RPS6KB2) and EIF4EBP1 (4E-BP1) by the mTORC1 signaling (PubMed:12271141, PubMed:24529379, PubMed:28215400, PubMed:33215753). The TSC-TBC complex is inactivated in response to nutrients, relieving inhibition of mTORC1 (PubMed:12172553, PubMed:24529379). Within the TSC-TBC complex, TSC1 stabilizes TSC2 and prevents TSC2 self-aggregation (PubMed:10585443, PubMed:28215400). Acts as a tumor suppressor (PubMed:9242607). Involved in microtubule-mediated protein transport via its ability to regulate mTORC1 signaling (By similarity). Also acts as a co-chaperone for HSP90AA1 facilitating HSP90AA1 chaperoning of protein clients such as kinases, TSC2 and glucocorticoid receptor NR3C1 (PubMed:29127155). Increases ATP binding to HSP90AA1 and inhibits HSP90AA1 ATPase activity (PubMed:29127155). Competes with the activating co-chaperone AHSA1 for binding to HSP90AA1, thereby providing a reciprocal regulatory mechanism for chaperoning of client proteins (PubMed:29127155). Recruits TSC2 to HSP90AA1 and stabilizes TSC2 by preventing the interaction between TSC2 and ubiquitin ligase HERC1 (PubMed:16464865, PubMed:29127155).

Cellular Location

Lysosome membrane; Peripheral membrane protein. Cytoplasm, cytosol Note=Recruited to lysosomal membranes in a RHEB-dependent process in absence of nutrients (PubMed:24529379). In response to nutrients, the complex dissociates from lysosomal membranes and relocates to the cytosol (PubMed:24529379).

Tissue Location

Highly expressed in skeletal muscle, followed by heart, brain, placenta, pancreas, lung, liver and kidney (PubMed:9242607). Also expressed in embryonic kidney cells (PubMed:9242607).

Goat Anti-Hamartin / TSC1 (isoform 1) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Goat Anti-Hamartin / TSC1 (isoform 1) Antibody - Images



AF1519a (0.1 $\mu\text{g/ml}$) staining of Human Brain lysate (35 μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



HEK293 overexpressing Human TSC1 (RC213332) and probed with AF1519a (mock transfection in first lane), tested by Origene.



AF1519a (3.8 µg/ml) staining of paraffin embedded Human Uterus. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

Goat Anti-Hamartin / TSC1 (isoform 1) Antibody - Background

This gene encodes a growth inhibitory protein thought to play a role in the stabilization of tuberin. Mutations in this gene have been associated with tuberous sclerosis. Alternative splicing results in multiple transcript variants.

Goat Anti-Hamartin / TSC1 (isoform 1) Antibody - References

Polymorphic variants in TSC1 and TSC2 and their association with breast cancer phenotypes. Mehta MS, et al. *Breast Cancer Res Treat*, 2010 Jul 25. PMID 20658316.
A Large-scale genetic association study of esophageal adenocarcinoma risk. Liu CY, et al. *Carcinogenesis*, 2010 Jul. PMID 20453000.
Tandem affinity purification and identification of the human TSC1 protein complex. Guo L, et al. *Acta Biochim Biophys Sin (Shanghai)*, 2010 Apr. PMID 20383465.
FOXO3a regulates glycolysis via transcriptional control of tumor suppressor TSC1. Khatri S, et al. *J Biol Chem*, 2010 May 21. PMID 20371605.
Expanding the tuberous sclerosis phenotype: mild disease caused by a TSC1 splicing mutation. Blyth M, et al. *J Neurol Neurosurg Psychiatry*, 2010 Mar. PMID 20185476.