

Goat Anti-Histamine receptor H1(internal) Antibody
Peptide-affinity purified goat antibody
Catalog # AF1530a

Specification

Goat Anti-Histamine receptor H1(internal) Antibody - Product Information

Application	WB
Primary Accession	P35367
Other Accession	NP_000852 , 3269
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	55784

Goat Anti-Histamine receptor H1(internal) Antibody - Additional Information

Gene ID [3269](#)

Other Names

Histamine H1 receptor, H1R, HH1R, HRH1

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-Histamine receptor H1(internal) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-Histamine receptor H1(internal) Antibody - Protein Information

Name HRH1 ([HGNC:5182](#))

Function

G-protein-coupled receptor for histamine, a biogenic amine that functions as an immune modulator and a neurotransmitter (PubMed: [33828102](http://www.uniprot.org/citations/33828102), PubMed: [8280179](http://www.uniprot.org/citations/8280179)). Through the H1 receptor, histamine mediates the contraction of smooth muscles and increases capillary permeability due to contraction of terminal venules. Also mediates neurotransmission in the central nervous system and thereby regulates circadian rhythms, emotional and locomotor activities as well as cognitive functions (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein

Goat Anti-Histamine receptor H1(internal) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-Histamine receptor H1(internal) Antibody - Images



AF1530a (0.5 µg/ml) staining of Human Heart lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Goat Anti-Histamine receptor H1(internal) Antibody - Background

Histamine is a ubiquitous messenger molecule released from mast cells, enterochromaffin-like cells, and neurons. Its various actions are mediated by histamine receptors H1, H2, H3 and H4. This gene was thought to be intronless until recently. The protein encoded by this gene is an integral membrane protein and belongs to the G protein-coupled receptor superfamily. It mediates the contraction of smooth muscles, the increase in capillary permeability due to contraction of terminal venules, the release of catecholamine from adrenal medulla, and neurotransmission in the central nervous system. Multiple alternatively spliced variants, encoding the same protein, have been identified.

Goat Anti-Histamine receptor H1(internal) Antibody - References

- Variation at the NFATC2 Locus Increases the Risk of Thiazolinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.
- Physiogenomic analysis of statin-treated patients: domain-specific counter effects within the ACACB gene on low-density lipoprotein cholesterol? Ruaño G, et al. Pharmacogenomics, 2010 Jul. PMID 20602615.
- New genetic associations detected in a host response study to hepatitis B vaccine. Davila S, et al.

Genes Immun, 2010 Apr. PMID 20237496.

Gene-centric association signals for lipids and apolipoproteins identified via the HumanCVD BeadChip. Talmud PJ, et al. Am J Hum Genet, 2009 Nov. PMID 19913121.

Histamine acting on H1 receptor promotes inhibition of proliferation via PLC, RAC, and JNK-dependent pathways. Notcovich C, et al. Exp Cell Res, 2010 Feb 1. PMID 19913013.