

**Goat Anti-Kcnj11 / Kir6.2 Antibody**  
Peptide-affinity purified goat antibody  
Catalog # AF1586a

**Specification**

---

**Goat Anti-Kcnj11 / Kir6.2 Antibody - Product Information**

Application	IHC, WB
Primary Accession	<a href="#">Q14654</a>
Other Accession	<a href="#">NP_034732</a> , <a href="#">3767</a> , <a href="#">16514 (mouse)</a> , <a href="#">83535 (rat)</a>
Reactivity	Human
Predicted	Mouse, Rat, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	43526

**Goat Anti-Kcnj11 / Kir6.2 Antibody - Additional Information**

**Gene ID** 3767

**Other Names**

ATP-sensitive inward rectifier potassium channel 11, IKATP, Inward rectifier K(+) channel Kir6.2, Potassium channel, inwardly rectifying subfamily J member 11, KCNJ11

**Format**

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat Anti-Kcnj11 / Kir6.2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat Anti-Kcnj11 / Kir6.2 Antibody - Protein Information**

**Name** KCNJ11

**Function**

This receptor is controlled by G proteins. Inward rectifier potassium channels are characterized by a greater tendency to allow potassium to flow into the cell rather than out of it. Their voltage dependence is regulated by the concentration of extracellular potassium; as external potassium is raised, the voltage range of the channel opening shifts to more positive voltages. The inward rectification is mainly due to the blockage of outward current by internal magnesium. Can be

blocked by extracellular barium (By similarity). Subunit of ATP-sensitive potassium channels (KATP). Can form cardiac and smooth muscle-type KATP channels with ABCC9. KCNJ11 forms the channel pore while ABCC9 is required for activation and regulation.

#### Cellular Location


Membrane; Multi-pass membrane protein.

#### Goat Anti-Kcnj11 / Kir6.2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Goat Anti-Kcnj11 / Kir6.2 Antibody - Images



AF1586a (3.8  $\mu\text{g/ml}$ ) staining of paraffin embedded Human Pancreas. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



AF1586a (0.01  $\mu\text{g/ml}$ ) staining of Human Muscle lysate (35  $\mu\text{g}$  protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

#### Goat Anti-Kcnj11 / Kir6.2 Antibody - References

Endoplasmic reticulum accumulation of Kir6.2 without activation of ER stress response in islet cells from adult Sur1 knockout mice. Marhfour I, et al. Cell Tissue Res, 2010 May. PMID 20383647.

Voltage-dependent gating in a voltage sensor-less ion channel. Kurata HT, et al. PLoS Biol, 2010 Feb 23. PMID 20208975.

Sarcolemmal ATP-sensitive K(+) channels control energy expenditure determining body weight. Alekseev AE, et al. Cell Metab, 2010 Jan. PMID 20074528.

Characterization and functional restoration of a potassium channel Kir6.2 pore mutation identified in congenital hyperinsulinism. Bushman JD, et al. J Biol Chem, 2010 Feb 26. PMID 20032456.

Impact of disease-causing SUR1 mutations on the KATP channel subunit interface probed with a rhodamine protection assay. Hosy E, et al. J Biol Chem, 2010 Jan 29. PMID 19933268.