

Goat Anti-LRRK2 / PARK8 (near C Terminus) Antibody
Peptide-affinity purified goat antibody
Catalog # AF1636a

Specification

Goat Anti-LRRK2 / PARK8 (near C Terminus) Antibody - Product Information

Application	IHC, IF
Primary Accession	Q5S007
Other Accession	NP_940980 , 120892
Reactivity	Human
Predicted	Cow
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	286103

Goat Anti-LRRK2 / PARK8 (near C Terminus) Antibody - Additional Information

Gene ID 120892

Other Names

Leucine-rich repeat serine/threonine-protein kinase 2, 2.7.11.1, Dardarin, LRRK2, PARK8

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-LRRK2 / PARK8 (near C Terminus) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-LRRK2 / PARK8 (near C Terminus) Antibody - Protein Information

Name LRRK2

Synonyms PARK8

Function

Serine/threonine-protein kinase which phosphorylates a broad range of proteins involved in multiple processes such as neuronal plasticity, innate immunity, autophagy, and vesicle trafficking (PubMed: [17114044](http://www.uniprot.org/citations/17114044)), PubMed: [20949042](http://www.uniprot.org/citations/20949042)),

PubMed: 21850687, PubMed: 22012985, PubMed: 23395371, PubMed: 24687852, PubMed: 25201882, PubMed: 26014385, PubMed: 26824392, PubMed: 27830463, PubMed: 28720718, PubMed: 29125462, PubMed: 29127255, PubMed: 29212815, PubMed: 30398148, PubMed: 30635421). Is a key regulator of RAB GTPases by regulating the GTP/GDP exchange and interaction partners of RABs through phosphorylation (PubMed: 26824392, PubMed: 28720718, PubMed: 29125462, PubMed: 29127255, PubMed: 29212815, PubMed: 30398148, PubMed: 30635421). Phosphorylates RAB3A, RAB3B, RAB3C, RAB3D, RAB5A, RAB5B, RAB5C, RAB8A, RAB8B, RAB10, RAB12, RAB35, and RAB43 (PubMed: 23395371, PubMed: 26824392, PubMed: 28720718, PubMed: 29125462, PubMed: 29127255, PubMed: 29212815, PubMed: 30398148, PubMed: 30635421). Regulates the RAB3IP-catalyzed GDP/GTP exchange for RAB8A through the phosphorylation of 'Thr-72' on RAB8A (PubMed: 26824392). Inhibits the interaction between RAB8A and GDI1 and/or GDI2 by phosphorylating 'Thr-72' on RAB8A (PubMed: 26824392). Regulates primary ciliogenesis through phosphorylation of RAB8A and RAB10, which promotes SHH signaling in the brain (PubMed: 29125462, PubMed: 30398148). Together with RAB29, plays a role in the retrograde trafficking pathway for recycling proteins, such as mannose-6-phosphate receptor (M6PR), between lysosomes and the Golgi apparatus in a retromer-dependent manner (PubMed: 23395371). Regulates neuronal process morphology in the intact central nervous system (CNS) (PubMed: 17114044). Plays a role in synaptic vesicle trafficking (PubMed: 24687852). Plays an important role in recruiting SEC16A to endoplasmic reticulum exit sites (ERES) and in regulating ER to Golgi vesicle-mediated transport and ERES organization (PubMed: 25201882). Positively regulates autophagy through a calcium-dependent activation of the CaMKK/AMPK signaling pathway (PubMed: 22012985). The process involves activation of nicotinic acid adenine dinucleotide phosphate (NAADP) receptors, increase in lysosomal pH, and calcium release from lysosomes (PubMed: 22012985). Phosphorylates PRDX3 (PubMed: 21850687

target="_blank">21850687). By phosphorylating APP on 'Thr-743', which promotes the production and the nuclear translocation of the APP intracellular domain (AICD), regulates dopaminergic neuron apoptosis (PubMed:28720718). Acts as a positive regulator of innate immunity by mediating phosphorylation of RIPK2 downstream of NOD1 and NOD2, thereby enhancing RIPK2 activation (PubMed:27830463). Independent of its kinase activity, inhibits the proteasomal degradation of MAPT, thus promoting MAPT oligomerization and secretion (PubMed:26014385). In addition, has GTPase activity via its Roc domain which regulates LRRK2 kinase activity (PubMed:18230735, PubMed:26824392, PubMed:28720718, PubMed:29125462, PubMed:29212815).

Cellular Location

Cytoplasmic vesicle. Perikaryon. Golgi apparatus membrane; Peripheral membrane protein. Cell projection, axon. Cell projection, dendrite. Endoplasmic reticulum membrane; Peripheral membrane protein. Cytoplasmic vesicle, secretory vesicle, synaptic vesicle membrane. Endosome {ECO:0000250|UniProtKB:Q5S006}. Lysosome Mitochondrion outer membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Note=Colocalized with RAB29 along tubular structures emerging from Golgi apparatus (PubMed:23395371). Localizes to endoplasmic reticulum exit sites (ERES), also known as transitional endoplasmic reticulum (tER) (PubMed:25201882).

Tissue Location


Expressed in pyramidal neurons in all cortical laminae of the visual cortex, in neurons of the substantia nigra pars compacta and caudate putamen (at protein level). Expressed in neutrophils (at protein level) (PubMed:29127255). Expressed in the brain. Expressed throughout the adult brain, but at a lower level than in heart and liver. Also expressed in placenta, lung, skeletal muscle, kidney and pancreas. In the brain, expressed in the cerebellum, cerebral cortex, medulla, spinal cord occipital pole, frontal lobe, temporal lobe and putamen. Expression is particularly high in brain dopaminergic areas.


Goat Anti-LRRK2 / PARK8 (near C Terminus) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-LRRK2 / PARK8 (near C Terminus) Antibody - Images






AF1636a (1.5 $\mu\text{g/ml}$) staining of paraffin embedded Human Hippocampus CA4. Microwaved antigen retrieval with citrate buffer pH 6, HRP-staining.



EB06550 Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 $\mu\text{g/ml}$) followed by Alexa Fluor 488 secondary antibody (2 $\mu\text{g/ml}$), showing vesicle staining. The nuclear stain is DAPI (blue).



Negative control: Unimmunized goat IgG (10 $\mu\text{g/ml}$) followed by Alexa Fluor 488 secondary antibody (2 $\mu\text{g/ml}$). EB06550 Immunofluorescence analysis of paraformaldehyde fixed A549 cells,

permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear and vesicle staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).

Goat Anti-LRRK2 / PARK8 (near C Terminus) Antibody - Background

This gene is a member of the leucine-rich repeat kinase family and encodes a protein with an ankryin repeat region, a leucine-rich repeat (LRR) domain, a kinase domain, a DFG-like motif, a RAS domain, a GTPase domain, a MLK-like domain, and a WD40 domain. The protein is present largely in the cytoplasm but also associates with the mitochondrial outer membrane. Mutations in this gene have been associated with Parkinson disease-8.

Goat Anti-LRRK2 / PARK8 (near C Terminus) Antibody - References

Olfactory heterogeneity in LRRK2 related Parkinsonism. Silveira-Moriyama L, et al. *Mov Disord*, 2010 Sep 3. PMID 20818658.

LRRK2 G2019S mutations are associated with an increased cancer risk in Parkinson disease. Saunders-Pullman R, et al. *Mov Disord*, 2010 Sep 3. PMID 20818610.

Low frequency of common LRRK2 mutations in Mexican patients with Parkinson's disease. Yescas P, et al. *Neurosci Lett*, 2010 Aug 18. PMID 20727385.

Absence of Commonly Reported Leucine-Rich Repeat Kinase 2 Mutations in Eastern Indian Parkinson's Disease Patients. Sanyal J, et al. *Genet Test Mol Biomarkers*, 2010 Aug 19. PMID 20722494.

Penetrance in Parkinson's disease related to the LRRK2 R1441G mutation in the Basque country (Spain). Ruiz-Martínez J, et al. *Mov Disord*, 2010 Aug 18. PMID 20721916.