

Goat Anti-PGAM1 / PGAM2 / PGAM4 Antibody
Peptide-affinity purified goat antibody
Catalog # AF1816a

Specification

Goat Anti-PGAM1 / PGAM2 / PGAM4 Antibody - Product Information

Application	WB
Primary Accession	P18669
Other Accession	NP_001025062 , 5223 , 5224 , 441531
Reactivity	Human
Predicted	Mouse, Rat, Pig, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	28804

Goat Anti-PGAM1 / PGAM2 / PGAM4 Antibody - Additional Information

Gene ID 5223

Other Names

Phosphoglycerate mutase 1, 3.1.3.13, 5.4.2.11, 5.4.2.4, BPG-dependent PGAM 1, Phosphoglycerate mutase isozyme B, PGAM-B, PGAM1, PGAMA

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-PGAM1 / PGAM2 / PGAM4 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-PGAM1 / PGAM2 / PGAM4 Antibody - Protein Information

Name PGAM1 ([HGNC:8888](#))

Synonyms PGAMA

Function

Catalyzes the interconversion of 2-phosphoglycerate and 3-phosphoglycerate a crucial step in glycolysis, by using 2,3-bisphosphoglycerate (PubMed:23653202). Also

catalyzes the interconversion of (2R)-2,3-bisphosphoglycerate and (2R)-3-phospho- glyceroyl phosphate (PubMed:23653202).

Tissue Location


Expressed in the liver and brain. Not found in the muscle.

Goat Anti-PGAM1 / PGAM2 / PGAM4 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-PGAM1 / PGAM2 / PGAM4 Antibody - Images



Antibody (0.03µg/ml) staining of Human Cerebellum (A) and Liver (B) lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



Antibody (0.03 μ g/ml) staining of Human Cerebellum (A) and Liver (B) lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Goat Anti-PGAM1 / PGAM2 / PGAM4 Antibody - Background

Phosphoglycerate mutase (PGAM) catalyzes the reversible reaction of 3-phosphoglycerate (3-PGA) to 2-phosphoglycerate (2-PGA) in the glycolytic pathway. The PGAM is a dimeric enzyme containing, in different tissues, different proportions of a slow-migrating muscle (MM) isozyme, a fast-migrating brain (BB) isozyme, and a hybrid form (MB). This gene encodes muscle-specific PGAM subunit. Mutations in this gene cause muscle phosphoglycerate mutase deficiency, also known as glycogen storage disease X.

Goat Anti-PGAM1 / PGAM2 / PGAM4 Antibody - References

Defining the human deubiquitinating enzyme interaction landscape. Sowa ME, et al. Cell, 2009 Jul 23. PMID 19615732.

The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334.

Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences. Strausberg RL, et al. Proc Natl Acad Sci U S A, 2002 Dec 24. PMID 12477932.

Manifesting heterozygotes in a Japanese family with a novel mutation in the muscle-specific phosphoglycerate mutase (PGAM-M) gene. Hadjigeorgiou GM, et al. Neuromuscul Disord, 1999 Oct. PMID 10545043.

The molecular genetic basis of muscle phosphoglycerate mutase (PGAM) deficiency. Tsujino S, et al. Am J Hum Genet, 1993 Mar. PMID 8447317.