

**Goat Anti-PINK1 Antibody**  
Peptide-affinity purified goat antibody  
Catalog # AF1830a

**Specification**

---

**Goat Anti-PINK1 Antibody - Product Information**

Application	WB, FC
Primary Accession	<a href="#">Q9BXM7</a>
Other Accession	<a href="#">NP_115785</a> , <a href="#">65018</a> , <a href="#">68943 (mouse)</a> , <a href="#">298575 (rat)</a>
Reactivity	Rat
Predicted	Human, Mouse
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	62769

**Goat Anti-PINK1 Antibody - Additional Information**

**Gene ID** 65018

**Other Names**

Serine/threonine-protein kinase PINK1, mitochondrial, 2.7.11.1, BRPK, PTEN-induced putative kinase protein 1, PINK1

**Format**

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat Anti-PINK1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat Anti-PINK1 Antibody - Protein Information**

**Name** PINK1

**Function**

Serine/threonine-protein kinase which protects against mitochondrial dysfunction during cellular stress by phosphorylating mitochondrial proteins such as PRKN and DNMI1L, to coordinate mitochondrial quality control mechanisms that remove and replace dysfunctional mitochondrial components (PubMed: <http://www.uniprot.org/citations/14607334>)

target="\_blank">14607334</a>, PubMed:<a href="http://www.uniprot.org/citations/15087508" target="\_blank">15087508</a>, PubMed:<a href="http://www.uniprot.org/citations/18443288" target="\_blank">18443288</a>, PubMed:<a href="http://www.uniprot.org/citations/18957282" target="\_blank">18957282</a>, PubMed:<a href="http://www.uniprot.org/citations/19229105" target="\_blank">19229105</a>, PubMed:<a href="http://www.uniprot.org/citations/19966284" target="\_blank">19966284</a>, PubMed:<a href="http://www.uniprot.org/citations/20404107" target="\_blank">20404107</a>, PubMed:<a href="http://www.uniprot.org/citations/20547144" target="\_blank">20547144</a>, PubMed:<a href="http://www.uniprot.org/citations/20798600" target="\_blank">20798600</a>, PubMed:<a href="http://www.uniprot.org/citations/22396657" target="\_blank">22396657</a>, PubMed:<a href="http://www.uniprot.org/citations/23620051" target="\_blank">23620051</a>, PubMed:<a href="http://www.uniprot.org/citations/23754282" target="\_blank">23754282</a>, PubMed:<a href="http://www.uniprot.org/citations/23933751" target="\_blank">23933751</a>, PubMed:<a href="http://www.uniprot.org/citations/24660806" target="\_blank">24660806</a>, PubMed:<a href="http://www.uniprot.org/citations/24751536" target="\_blank">24751536</a>, PubMed:<a href="http://www.uniprot.org/citations/24784582" target="\_blank">24784582</a>, PubMed:<a href="http://www.uniprot.org/citations/24896179" target="\_blank">24896179</a>, PubMed:<a href="http://www.uniprot.org/citations/24898855" target="\_blank">24898855</a>, PubMed:<a href="http://www.uniprot.org/citations/25527291" target="\_blank">25527291</a>, PubMed:<a href="http://www.uniprot.org/citations/32484300" target="\_blank">32484300</a>). Depending on the severity of mitochondrial damage and/or dysfunction, activity ranges from preventing apoptosis and stimulating mitochondrial biogenesis to regulating mitochondrial dynamics and eliminating severely damaged mitochondria via mitophagy (PubMed:<a href="http://www.uniprot.org/citations/15087508" target="\_blank">15087508</a>, PubMed:<a href="http://www.uniprot.org/citations/18443288" target="\_blank">18443288</a>, PubMed:<a href="http://www.uniprot.org/citations/19966284" target="\_blank">19966284</a>, PubMed:<a href="http://www.uniprot.org/citations/20404107" target="\_blank">20404107</a>, PubMed:<a href="http://www.uniprot.org/citations/20798600" target="\_blank">20798600</a>, PubMed:<a href="http://www.uniprot.org/citations/22396657" target="\_blank">22396657</a>, PubMed:<a href="http://www.uniprot.org/citations/23620051" target="\_blank">23620051</a>, PubMed:<a href="http://www.uniprot.org/citations/24898855" target="\_blank">24898855</a>, PubMed:<a href="http://www.uniprot.org/citations/32047033" target="\_blank">32047033</a>, PubMed:<a href="http://www.uniprot.org/citations/32484300" target="\_blank">32484300</a>). Mediates the translocation and activation of PRKN at the outer membrane (OMM) of dysfunctional/depolarized mitochondria (PubMed:<a href="http://www.uniprot.org/citations/19966284" target="\_blank">19966284</a>, PubMed:<a href="http://www.uniprot.org/citations/20404107" target="\_blank">20404107</a>, PubMed:<a href="http://www.uniprot.org/citations/20798600" target="\_blank">20798600</a>, PubMed:<a href="http://www.uniprot.org/citations/23754282" target="\_blank">23754282</a>, PubMed:<a href="http://www.uniprot.org/citations/24660806" target="\_blank">24660806</a>, PubMed:<a href="http://www.uniprot.org/citations/24751536" target="\_blank">24751536</a>, PubMed:<a href="http://www.uniprot.org/citations/24784582" target="\_blank">24784582</a>, PubMed:<a href="http://www.uniprot.org/citations/25474007" target="\_blank">25474007</a>, PubMed:<a href="http://www.uniprot.org/citations/25527291" target="\_blank">25527291</a>). At the OMM of damaged mitochondria, phosphorylates pre-existing polyubiquitin chains at 'Ser-65', the PINK1-phosphorylated polyubiquitin then recruits PRKN from the cytosol to the OMM where PRKN is fully activated by phosphorylation at 'Ser-65' by PINK1 (PubMed:<a href="http://www.uniprot.org/citations/19966284" target="\_blank">19966284</a>, PubMed:<a href="http://www.uniprot.org/citations/20404107" target="\_blank">20404107</a>, PubMed:<a href="http://www.uniprot.org/citations/20798600" target="\_blank">20798600</a>, PubMed:<a href="http://www.uniprot.org/citations/23754282" target="\_blank">23754282</a>, PubMed:<a href="http://www.uniprot.org/citations/24660806" target="\_blank">24660806</a>, PubMed:<a href="http://www.uniprot.org/citations/24751536" target="\_blank">24751536</a>, PubMed:<a href="http://www.uniprot.org/citations/24784582" target="\_blank">24784582</a>, PubMed:<a href="http://www.uniprot.org/citations/25474007" target="\_blank">25474007</a>, PubMed:<a href="http://www.uniprot.org/citations/25527291" target="\_blank">25527291</a>). In damaged mitochondria, mediates the decision between mitophagy or preventing apoptosis by promoting PRKN-dependent poly- or monoubiquitination of VDAC1; polyubiquitination of VDAC1 by PRKN

promotes mitophagy, while monoubiquitination of VDAC1 by PRKN decreases mitochondrial calcium influx which ultimately inhibits apoptosis (PubMed:<a href="http://www.uniprot.org/citations/32047033" target="\_blank">32047033</a>). When cellular stress results in irreversible mitochondrial damage, functions with PRKN to promote clearance of damaged mitochondria via selective autophagy (mitophagy) (PubMed:<a href="http://www.uniprot.org/citations/14607334" target="\_blank">14607334</a>, PubMed:<a href="http://www.uniprot.org/citations/15087508" target="\_blank">15087508</a>, PubMed:<a href="http://www.uniprot.org/citations/19966284" target="\_blank">19966284</a>, PubMed:<a href="http://www.uniprot.org/citations/20404107" target="\_blank">20404107</a>, PubMed:<a href="http://www.uniprot.org/citations/20798600" target="\_blank">20798600</a>, PubMed:<a href="http://www.uniprot.org/citations/23933751" target="\_blank">23933751</a>). The PINK1-PRKN pathway also promotes fission of damaged mitochondria by phosphorylating and thus promoting the PRKN-dependent degradation of mitochondrial proteins involved in fission such as MFN2 (PubMed:<a href="http://www.uniprot.org/citations/18443288" target="\_blank">18443288</a>, PubMed:<a href="http://www.uniprot.org/citations/23620051" target="\_blank">23620051</a>, PubMed:<a href="http://www.uniprot.org/citations/24898855" target="\_blank">24898855</a>). This prevents the refusion of unhealthy mitochondria with the mitochondrial network or initiates mitochondrial fragmentation facilitating their later engulfment by autophagosomes (PubMed:<a href="http://www.uniprot.org/citations/18443288" target="\_blank">18443288</a>, PubMed:<a href="http://www.uniprot.org/citations/23620051" target="\_blank">23620051</a>). Also promotes mitochondrial fission independently of PRKN and ATG7-mediated mitophagy, via the phosphorylation and activation of DNM1L (PubMed:<a href="http://www.uniprot.org/citations/18443288" target="\_blank">18443288</a>, PubMed:<a href="http://www.uniprot.org/citations/32484300" target="\_blank">32484300</a>). Regulates motility of damaged mitochondria by promoting the ubiquitination and subsequent degradation of MIRO1 and MIRO2; in motor neurons, this likely inhibits mitochondrial intracellular anterograde transport along the axons which probably increases the chance of the mitochondria undergoing mitophagy in the soma (PubMed:<a href="http://www.uniprot.org/citations/22396657" target="\_blank">22396657</a>). Required for ubiquinone reduction by mitochondrial complex I by mediating phosphorylation of complex I subunit NDUFA10 (By similarity). Phosphorylates LETM1, positively regulating its mitochondrial calcium transport activity (PubMed:<a href="http://www.uniprot.org/citations/29123128" target="\_blank">29123128</a>).

### Cellular Location

Mitochondrion outer membrane; Single-pass membrane protein. Mitochondrion inner membrane {ECO:0000250|UniProtKB:Q99MQ3}; Single-pass membrane protein. Cytoplasm, cytosol. Note=Localizes mostly in mitochondrion and the two smaller proteolytic processed fragments localize mainly in cytosol (PubMed:19229105). When mitochondria lose mitochondrial membrane potential following damage, PINK1 import is arrested, which induces its accumulation in the outer mitochondrial membrane, where it acquires kinase activity (PubMed:18957282)

### Tissue Location

Highly expressed in heart, skeletal muscle and testis, and at lower levels in brain, placenta, liver, kidney, pancreas, prostate, ovary and small intestine. Present in the embryonic testis from an early stage of development


## Goat Anti-PINK1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Goat Anti-PINK1 Antibody - Images



AF1830a (1  $\mu$ g/ml) staining of Rat Testis lysate (35  $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB07940 Flow cytometric analysis of paraformaldehyde fixed Jurkat cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.

### Goat Anti-PINK1 Antibody - Background

This gene encodes a serine/threonine protein kinase that localizes to mitochondria. It is thought to protect cells from stress-induced mitochondrial dysfunction. Mutations in this gene cause one form of autosomal recessive early-onset Parkinson disease.

### Goat Anti-PINK1 Antibody - References

Oxidative stress alters the regulatory control of p66Shc and Akt in PINK1 deficient cells. Maj MC, et al. *Biochem Biophys Res Commun*, 2010 Aug 27. PMID 20637729.  
Assessing the prevalence of PINK1 genetic variants in South African patients diagnosed with early- and late-onset Parkinson's disease. Keyser RJ, et al. *Biochem Biophys Res Commun*, 2010 Jul 16.

PMID 20558144.

Progression of subtle motor signs in PINK1 mutation carriers with mild dopaminergic deficit. Eggers C, et al. *Neurology*, 2010 Jun 1. PMID 20513816.

Structural imaging in the presymptomatic stage of genetically determined parkinsonism. Reetz K, et al. *Neurobiol Dis*, 2010 Sep. PMID 20483373.

Clinical and demographic characteristics of PINK1 mutation carriers--a meta-analysis. Kasten M, et al. *Mov Disord*, 2010 May 15. PMID 20461815.