

**Goat Anti-PRDX1 Antibody**  
Peptide-affinity purified goat antibody  
Catalog # AF1863a

## Specification

---

### Goat Anti-PRDX1 Antibody - Product Information

Application	WB, IHC
Primary Accession	<a href="#">Q06830</a>
Other Accession	<a href="#">NP_859048</a> , <a href="#">5052</a> , <a href="#">18477 (mouse)</a> , <a href="#">117254 (rat)</a>
Reactivity	Human
Predicted	Mouse, Rat, Pig, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	22110

### Goat Anti-PRDX1 Antibody - Additional Information

**Gene ID** 5052

#### Other Names

Peroxiredoxin-1, 1.11.1.15, Natural killer cell-enhancing factor A, NKEF-A, Proliferation-associated gene protein, PAG, Thioredoxin peroxidase 2, Thioredoxin-dependent peroxide reductase 2, PRDX1, PAGA, PAGB, TDPX2

#### Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

Goat Anti-PRDX1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### Goat Anti-PRDX1 Antibody - Protein Information

**Name** PRDX1

**Synonyms** PAGA, PAGB, TDPX2

#### Function

Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic hydroperoxides to water and alcohols, respectively. Plays a role in cell protection against oxidative

stress by detoxifying peroxides and as sensor of hydrogen peroxide-mediated signaling events. Might participate in the signaling cascades of growth factors and tumor necrosis factor-alpha by regulating the intracellular concentrations of H<sub>2</sub>O<sub>2</sub> (PubMed:<a href="http://www.uniprot.org/citations/9497357" target="\_blank">9497357</a>). Reduces an intramolecular disulfide bond in GDPD5 that gates the ability to GDPD5 to drive postmitotic motor neuron differentiation (By similarity).

#### Cellular Location

Cytoplasm. Melanosome Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV

#### Goat Anti-PRDX1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


#### Goat Anti-PRDX1 Antibody - Images



AF1863a (0.01 µg/ml) staining of HepG2 lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



AF1863a (2.5  $\mu\text{g/ml}$ ) staining of paraffin embedded Human Thyroid Gland. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



AF1863a (2.5  $\mu\text{g/ml}$ ) staining of paraffin embedded Human Placenta. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.