

**Goat Anti-PSPH Antibody**  
Peptide-affinity purified goat antibody  
Catalog # AF1876a

## Specification

---

### Goat Anti-PSPH Antibody - Product Information

Application	<b>WB</b>
Primary Accession	<a href="#">P78330</a>
Other Accession	<a href="#">NP_004568</a> , <a href="#">5723</a> , <a href="#">100678 (mouse)</a> , <a href="#">304429 (rat)</a>
Reactivity	<b>Human, Mouse</b>
Predicted	<b>Rat</b>
Host	<b>Goat</b>
Clonality	<b>Polyclonal</b>
Concentration	<b>100ug/200ul</b>
Isotype	<b>IgG</b>
Calculated MW	<b>25008</b>

### Goat Anti-PSPH Antibody - Additional Information

**Gene ID** 5723

#### Other Names

Phosphoserine phosphatase, PSP, PSPase, 3.1.3.3, L-3-phosphoserine phosphatase, O-phosphoserine phosphohydrolase, PSPH

#### Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

Goat Anti-PSPH Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### Goat Anti-PSPH Antibody - Protein Information

**Name** PSPH ([HGNC:9577](#))

#### Function

Catalyzes the last irreversible step in the biosynthesis of L-serine from carbohydrates, the dephosphorylation of O-phospho-L-serine to L-serine (PubMed:[12213811](http://www.uniprot.org/citations/12213811), PubMed:[14673469](http://www.uniprot.org/citations/14673469), PubMed:[14673469](http://www.uniprot.org/citations/14673469), PubMed:[14673469](http://www.uniprot.org/citations/14673469))

[15291819](http://www.uniprot.org/citations/15291819), PubMed: [25080166](http://www.uniprot.org/citations/25080166), PubMed: [9222972](http://www.uniprot.org/citations/9222972)). L-serine can then be used in protein synthesis, to produce other amino acids, in nucleotide metabolism or in glutathione synthesis, or can be racemized to D-serine, a neuromodulator (PubMed: [14673469](http://www.uniprot.org/citations/14673469)). May also act on O-phospho-D-serine (Probable).

#### Cellular Location

Cytoplasm, cytosol.

#### Goat Anti-PSPH Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Goat Anti-PSPH Antibody - Images



AF1876a (0.05 µg/ml) staining of Mouse Brain lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

#### Goat Anti-PSPH Antibody - Background

The protein encoded by this gene belongs to a subfamily of the phosphotransferases. This encoded enzyme is responsible for the third and last step in L-serine formation. It catalyzes magnesium-dependent hydrolysis of L-phosphoserine and is also involved in an exchange reaction between L-serine and L-phosphoserine. Deficiency of this protein is thought to be linked to Williams syndrome.

#### Goat Anti-PSPH Antibody - References

Large-scale mapping of human protein-protein interactions by mass spectrometry. Ewing RM, et al. Mol Syst Biol, 2007. PMID 17353931.  
The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene

Collection (MGC). Gerhard DS, et al. Genome Res, 2004 Oct. PMID 15489334.

How calcium inhibits the magnesium-dependent enzyme human phosphoserine phosphatase.

Peeraer Y, et al. Eur J Biochem, 2004 Aug. PMID 15291819.

Complete sequencing and characterization of 21,243 full-length human cDNAs. Ota T, et al. Nat Genet, 2004 Jan. PMID 14702039.

Mutations responsible for 3-phosphoserine phosphatase deficiency. Veiga-da-Cunha M, et al. Eur J Hum Genet, 2004 Feb. PMID 14673469.