

Goat Anti-ribosomal protein L8 Antibody
Peptide-affinity purified goat antibody
Catalog # AF1928a

Specification

Goat Anti-ribosomal protein L8 Antibody - Product Information

Application	WB, IHC
Primary Accession	P62917
Other Accession	NP_150644 , 6132 , 26961 (mouse) , 26962 (rat)
Reactivity	Human
Predicted	Mouse, Rat, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	28025

Goat Anti-ribosomal protein L8 Antibody - Additional Information

Gene ID 6132

Other Names

60S ribosomal protein L8, RPL8

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-ribosomal protein L8 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-ribosomal protein L8 Antibody - Protein Information

Name RPL8

Function

Component of the large ribosomal subunit. The ribosome is a large ribonucleoprotein complex responsible for the synthesis of proteins in the cell.

Cellular Location

Cytoplasm.

Goat Anti-ribosomal protein L8 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-ribosomal protein L8 Antibody - Images



AF1928a (0.01 $\mu\text{g/ml}$) staining of HepG2 cell lysate (35 μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



AF1928a (2.5 $\mu\text{g/ml}$) staining of paraffin embedded Human Cerebral Cortex. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.

Goat Anti-ribosomal protein L8 Antibody - Background

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L2P family of ribosomal proteins. It is located in the cytoplasm. In rat, the protein associates with the 5.8S rRNA, very likely participates in the binding of aminoacyl-tRNA, and is a constituent of the elongation factor 2-binding site at the ribosomal subunit interface. Alternatively spliced transcript variants encoding the same protein exist. As is typical for

genes encoding ribosomal proteins, there are multiple processed pseudogenes of this gene dispersed through the genome.

Goat Anti-ribosomal protein L8 Antibody - References

Molecular characterization of the response to chemotherapy in conventional osteosarcomas: predictive value of HSD17B10 and IFITM2. Salas S, et al. *Int J Cancer*, 2009 Aug 15. PMID 19449377.

A human protein-protein interaction network: a resource for annotating the proteome. Stelzl U, et al. *Cell*, 2005 Sep 23. PMID 16169070.

Nucleolar proteome dynamics. Andersen JS, et al. *Nature*, 2005 Jan 6. PMID 15635413.

Immunoaffinity profiling of tyrosine phosphorylation in cancer cells. Rush J, et al. *Nat Biotechnol*, 2005 Jan. PMID 15592455.

The status, quality, and expansion of the NIH full-length cDNA project: the Mammalian Gene Collection (MGC). Gerhard DS, et al. *Genome Res*, 2004 Oct. PMID 15489334.