

**Goat Anti-RXR alpha Antibody**  
Peptide-affinity purified goat antibody  
Catalog # AF1947a

## Specification

---

### Goat Anti-RXR alpha Antibody - Product Information

Application	WB
Primary Accession	<a href="#">P19793</a>
Other Accession	<a href="#">NP_002948</a> , <a href="#">6256</a>
Reactivity	Human
Predicted	Mouse, Rat
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	50811

### Goat Anti-RXR alpha Antibody - Additional Information

**Gene ID** 6256

#### Other Names

Retinoic acid receptor RXR-alpha, Nuclear receptor subfamily 2 group B member 1, Retinoid X receptor alpha, RXRA, NR2B1

#### Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

Goat Anti-RXR alpha Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### Goat Anti-RXR alpha Antibody - Protein Information

**Name** RXRA

**Synonyms** NR2B1

#### Function

Receptor for retinoic acid that acts as a transcription factor (PubMed:<a href="http://www.uniprot.org/citations/11162439" target="\_blank">11162439</a>, PubMed:<a href="http://www.uniprot.org/citations/11915042" target="\_blank">11915042</a>, PubMed:<a href="http://www.uniprot.org/citations/11915042" target="\_blank">11915042</a>, PubMed:<a href="http://www.uniprot.org/citations/11915042" target="\_blank">11915042</a>)

href="http://www.uniprot.org/citations/37478846" target="\_blank">37478846</a>). Forms homo- or heterodimers with retinoic acid receptors (RARs) and binds to target response elements in response to their ligands, all-trans or 9- cis retinoic acid, to regulate gene expression in various biological processes (PubMed:<a href="http://www.uniprot.org/citations/10195690" target="\_blank">10195690</a>, PubMed:<a href="http://www.uniprot.org/citations/11162439" target="\_blank">11162439</a>, PubMed:<a href="http://www.uniprot.org/citations/11915042" target="\_blank">11915042</a>, PubMed:<a href="http://www.uniprot.org/citations/16107141" target="\_blank">16107141</a>, PubMed:<a href="http://www.uniprot.org/citations/17761950" target="\_blank">17761950</a>, PubMed:<a href="http://www.uniprot.org/citations/18800767" target="\_blank">18800767</a>, PubMed:<a href="http://www.uniprot.org/citations/19167885" target="\_blank">19167885</a>, PubMed:<a href="http://www.uniprot.org/citations/28167758" target="\_blank">28167758</a>, PubMed:<a href="http://www.uniprot.org/citations/37478846" target="\_blank">37478846</a>). The RAR/RXR heterodimers bind to the retinoic acid response elements (RARE) composed of tandem 5'-AGGTCA-3' sites known as DR1-DR5 to regulate transcription (PubMed:<a href="http://www.uniprot.org/citations/10195690" target="\_blank">10195690</a>, PubMed:<a href="http://www.uniprot.org/citations/11162439" target="\_blank">11162439</a>, PubMed:<a href="http://www.uniprot.org/citations/11915042" target="\_blank">11915042</a>, PubMed:<a href="http://www.uniprot.org/citations/17761950" target="\_blank">17761950</a>, PubMed:<a href="http://www.uniprot.org/citations/28167758" target="\_blank">28167758</a>). The high affinity ligand for retinoid X receptors (RXRs) is 9-cis retinoic acid (PubMed:<a href="http://www.uniprot.org/citations/1310260" target="\_blank">1310260</a>). In the absence of ligand, the RXR-RAR heterodimers associate with a multiprotein complex containing transcription corepressors that induce histone deacetylation, chromatin condensation and transcriptional suppression (PubMed:<a href="http://www.uniprot.org/citations/20215566" target="\_blank">20215566</a>). On ligand binding, the corepressors dissociate from the receptors and coactivators are recruited leading to transcriptional activation (PubMed:<a href="http://www.uniprot.org/citations/20215566" target="\_blank">20215566</a>, PubMed:<a href="http://www.uniprot.org/citations/37478846" target="\_blank">37478846</a>, PubMed:<a href="http://www.uniprot.org/citations/9267036" target="\_blank">9267036</a>). Serves as a common heterodimeric partner for a number of nuclear receptors, such as RARA, RARB and PPARA (PubMed:<a href="http://www.uniprot.org/citations/10195690" target="\_blank">10195690</a>, PubMed:<a href="http://www.uniprot.org/citations/11915042" target="\_blank">11915042</a>, PubMed:<a href="http://www.uniprot.org/citations/28167758" target="\_blank">28167758</a>, PubMed:<a href="http://www.uniprot.org/citations/29021580" target="\_blank">29021580</a>). The RXRA/RARB heterodimer can act as a transcriptional repressor or transcriptional activator, depending on the RARE DNA element context (PubMed:<a href="http://www.uniprot.org/citations/29021580" target="\_blank">29021580</a>). The RXRA/PPARA heterodimer is required for PPARA transcriptional activity on fatty acid oxidation genes such as ACOX1 and the P450 system genes (PubMed:<a href="http://www.uniprot.org/citations/10195690" target="\_blank">10195690</a>). Together with RARA, positively regulates microRNA- 10a expression, thereby inhibiting the GATA6/VCAM1 signaling response to pulsatile shear stress in vascular endothelial cells (PubMed:<a href="http://www.uniprot.org/citations/28167758" target="\_blank">28167758</a>). Acts as an enhancer of RARA binding to RARE DNA element (PubMed:<a href="http://www.uniprot.org/citations/28167758" target="\_blank">28167758</a>). May facilitate the nuclear import of heterodimerization partners such as VDR and NR4A1 (PubMed:<a href="http://www.uniprot.org/citations/12145331" target="\_blank">12145331</a>, PubMed:<a href="http://www.uniprot.org/citations/15509776" target="\_blank">15509776</a>). Promotes myelin debris phagocytosis and remyelination by macrophages (PubMed:<a href="http://www.uniprot.org/citations/26463675" target="\_blank">26463675</a>). Plays a role in the attenuation of the innate immune system in response to viral infections, possibly by negatively regulating the transcription of antiviral genes such as type I IFN genes (PubMed:<a href="http://www.uniprot.org/citations/25417649" target="\_blank">25417649</a>). Involved in the regulation of calcium signaling by repressing ITPR2 gene expression, thereby controlling cellular senescence (PubMed:<a href="http://www.uniprot.org/citations/30216632" target="\_blank">30216632</a>).

### Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00407, ECO:0000269|PubMed:11915042, ECO:0000269|PubMed:12145331, ECO:0000269|PubMed:15509776, ECO:0000269|PubMed:17761950, ECO:0000269|PubMed:28167758}. Cytoplasm. Mitochondrion. Note=Localization to the nucleus is enhanced by vitamin D3 (PubMed:15509776). Nuclear localization may be enhanced by the interaction with heterodimerization partner VDR (PubMed:12145331). Translocation to the mitochondrion upon interaction with NR4A1 (PubMed:15509776, PubMed:17761950). Increased nuclear localization upon pulsatile shear stress (PubMed:28167758)

### Tissue Location


Expressed in lung fibroblasts (at protein level) (PubMed:30216632). Expressed in monocytes (PubMed:26463675). Highly expressed in liver, also found in kidney and brain (PubMed:14702039, PubMed:2159111, PubMed:24275569).

### Goat Anti-RXR alpha Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Goat Anti-RXR alpha Antibody - Images



AF1947a (0.5 µg/ml) staining of A431 lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

### Goat Anti-RXR alpha Antibody - Background

Retinoid X receptors (RXRs) and retinoic acid receptors (RARs), are nuclear receptors that mediate the biological effects of retinoids by their involvement in retinoic acid-mediated gene activation. These receptors exert their action by binding, as homodimers or heterodimers, to specific sequences in the promoters of target genes and regulating their transcription. The protein encoded by this gene is a member of the steroid and thyroid hormone receptor superfamily of transcriptional

regulators.

### **Goat Anti-RXR alpha Antibody - References**

- Variation at the NFATC2 Locus Increases the Risk of Thiazolidinedione-Induced Edema in the Diabetes REduction Assessment with ramipril and rosiglitazone Medication (DREAM) Study. Bailey SD, et al. Diabetes Care, 2010 Jul 13. PMID 20628086.
- Physiogenomic analysis of statin-treated patients: domain-specific counter effects within the ACACB gene on low-density lipoprotein cholesterol? Ruaño G, et al. Pharmacogenomics, 2010 Jul. PMID 20602615.
- Genetic variation in the retinoid X receptor and calcium-sensing receptor and risk of colorectal cancer in the Colon Cancer Family Registry. Jacobs ET, et al. Carcinogenesis, 2010 Aug. PMID 20558521.
- NSAID sulindac and its analog bind RXRalpha and inhibit RXRalpha-dependent AKT signaling. Zhou H, et al. Cancer Cell, 2010 Jun 15. PMID 20541701.
- Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614.