

Goat Anti-RXR gamma Antibody
Peptide-affinity purified goat antibody
Catalog # AF1949a**Specification**

Goat Anti-RXR gamma Antibody - Product Information

Application	WB
Primary Accession	P48443
Other Accession	NP_008848 , 6258
Reactivity	Human
Predicted	Mouse, Rat, Pig, Dog
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	50871

Goat Anti-RXR gamma Antibody - Additional Information**Gene ID** 6258**Other Names**

Retinoic acid receptor RXR-gamma, Nuclear receptor subfamily 2 group B member 3, Retinoid X receptor gamma, RXRG, NR2B3

Format

0.5 mg IgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-RXR gamma Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-RXR gamma Antibody - Protein Information**Name** RXRG**Synonyms** NR2B3**Function**

Receptor for retinoic acid. Retinoic acid receptors bind as heterodimers to their target response elements in response to their ligands, all-trans or 9-cis retinoic acid, and regulate gene expression in various biological processes. The RAR/RXR heterodimers bind to the retinoic acid response

elements (RARE) composed of tandem 5'-AGGTCA-3' sites known as DR1-DR5. The high affinity ligand for RXRs is 9-cis retinoic acid (By similarity).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00407, ECO:0000269|PubMed:28167758}.
Cytoplasm

Tissue Location


Expressed in aortic endothelial cells (at protein level).

Goat Anti-RXR gamma Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-RXR gamma Antibody - Images



AF1949a (1 µg/ml) staining of HeLa lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Goat Anti-RXR gamma Antibody - Background

This gene encodes a member of the retinoid X receptor (RXR) family of nuclear receptors which are involved in mediating the antiproliferative effects of retinoic acid (RA). This receptor forms dimers with the retinoic acid, thyroid hormone, and vitamin D receptors, increasing both DNA binding and transcriptional function on their respective response elements. This gene is expressed at significantly lower levels in non-small cell lung cancer cells. Alternatively spliced transcript variants have been described.

Goat Anti-RXR gamma Antibody - References

Maternal genes and facial clefts in offspring: a comprehensive search for genetic associations in two population-based cleft studies from Scandinavia. Jugessur A, et al. PLoS One, 2010 Jul 9. PMID 20634891.

Physiogenomic analysis of statin-treated patients: domain-specific counter effects within the ACACB gene on low-density lipoprotein cholesterol? Ruaño G, et al. Pharmacogenomics, 2010 Jul. PMID 20602615.

Personalized smoking cessation: interactions between nicotine dose, dependence and quit-success genotype score. Rose JE, et al. Mol Med, 2010 Jul-Aug. PMID 20379614.

LIM protein Ajuba functions as a nuclear receptor corepressor and negatively regulates retinoic acid signaling. Hou Z, et al. Proc Natl Acad Sci U S A, 2010 Feb 16. PMID 20133701.

Epigenetic inactivation of retinoid X receptor genes in non-small cell lung cancer and the relationship with clinicopathologic features. Lee SM, et al. Cancer Genet Cytogenet, 2010 Feb. PMID 20113835.