

Goat Anti-SAR1B / SARA2 Antibody

Peptide-affinity purified goat antibody Catalog # AF1958a

Specification

Goat Anti-SAR1B / SARA2 Antibody - Product Information

Application Primary Accession Other Accession

Reactivity Predicted Host Clonality Concentration Isotype Calculated MW WB, E <u>Q9Y6B6</u> <u>NP_057187</u>, <u>51128</u>, <u>66397 (mouse)</u>, <u>287276</u> (<u>rat</u>) Mouse, Rat Human Goat Polyclonal 100ug/200ul IgG 22410

Goat Anti-SAR1B / SARA2 Antibody - Additional Information

Gene ID 51128

Other Names GTP-binding protein SAR1b, GTP-binding protein B, GTBPB, SAR1B, SARA2, SARB

Dilution WB~~1:1000 E~~N/A

Format

0.5 mg lgG/ml in Tris saline (20mM Tris pH7.3, 150mM NaCl), 0.02% sodium azide, with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions Goat Anti-SAR1B / SARA2 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-SAR1B / SARA2 Antibody - Protein Information

Name SAR1B {ECO:0000303|PubMed:33186557, ECO:0000312|HGNC:HGNC:10535}

Function

Small GTPase that cycles between an active GTP-bound and an inactive GDP-bound state and



mainly functions in vesicle-mediated endoplasmic reticulum (ER) to Golgi transport. The active GTP-bound form inserts into the endoplasmic reticulum membrane where it recruits the remainder of the coat protein complex II/COPII (PubMed:23433038, PubMed:23433038, PubMed:33386557, PubMed:33186557, PubMed:33186557, PubMed:36369712). The coat protein complex II assembling and polymerizing on endoplasmic reticulum membrane is responsible for both the sorting of cargos and the deformation and budding of membranes into vesicles destined to the Golgi (PubMed:23433038, PubMed:23433038, PubMed:23433038, PubMed:23433038, PubMed:23433038, PubMed:33186557, NubMed:33186557</a

href="http://www.uniprot.org/citations/32358066" target="_blank">32358066, PubMed:33186557). In addition to its role in vesicle trafficking, can also function as a leucine sensor regulating TORC1 signaling and more indirectly cellular metabolism, growth and survival. In absence of leucine, interacts with the GATOR2 complex via MIOS and inhibits TORC1 signaling. The binding of leucine abrogates the interaction with GATOR2 and the inhibition of the TORC1 signaling. This function is completely independent of the GTPase activity of SAR1B (PubMed:34290409).

Cellular Location

Endoplasmic reticulum membrane; Peripheral membrane protein

{ECO:0000250|UniProtKB:Q9QVY3}. Golgi apparatus, Golgi stack membrane

{ECO:0000250|UniProtKB:Q9QVY3}; Peripheral membrane protein

{ECO:0000250|UniProtKB:Q9QVY3}. Cytoplasm, cytosol. Lysosome membrane. Note=Active at endoplasmic reticulum exit sites (ERES) where it inserts into the membrane and recruits the remainder of the coat protein complex II/COPII (PubMed:23433038). Upon leucine deprivation, associates with lysosomal membranes to repress TORC1 signaling (PubMed:34290409).

Tissue Location

Expressed in many tissues including small intestine, liver, muscle and brain.

Goat Anti-SAR1B / SARA2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Goat Anti-SAR1B / SARA2 Antibody - Images



250kDa 150kDa 100kDa 75kDa 30kDa 37kDa 25kDa 20kDa

15kDa

AF1958a (0.03 μ g/ml) staining of Mouse Liver lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

HEK293 overexpressing SAR1B (RC210593) and probed with AF1958a (mock transfection in first lane), tested by Origene.

Goat Anti-SAR1B / SARA2 Antibody - Background

The protein encoded by this gene is a small GTPase that acts as a homodimer. The encoded protein is activated by the guanine nucleotide exchange factor PREB and is involved in protein transport from the endoplasmic reticulum to the Golgi. This protein is part of the COPII coat complex. Defects in this gene are a cause of chylomicron retention disease (CMRD), also known as Anderson disease (ANDD). Two transcript variants encoding the same protein have been found for this gene.

Goat Anti-SAR1B / SARA2 Antibody - References

Variable phenotypic expression of chylomicron retention disease in a kindred carrying a mutation of the Sara2 gene. Cefal AB, et al. Metabolism, 2010 Apr. PMID 19846172.

Anderson's disease (chylomicron retention disease): a new mutation in the SARA2 gene associated with muscular and cardiac abnormalities. Silvain M, et al. Clin Genet, 2008 Dec. PMID 18786134. Toward a confocal subcellular atlas of the human proteome. Barbe L, et al. Mol Cell Proteomics, 2008 Mar. PMID 18029348.

Anderson or chylomicron retention disease: molecular impact of five mutations in the SAR1B gene on the structure and the functionality of Sar1b protein. Charcosset M, et al. Mol Genet Metab, 2008 Jan. PMID 17945526.

Expression of Sara2 human gene in erythroid progenitors. Jardim DL, et al. J Biochem Mol Biol, 2005



May 31. PMID 15943909.