

PARP12 Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF2498a

Specification

PARP12 Antibody (internal region) - Product Information

Application	IF, E
Primary Accession	O9H0J9
Other Accession	NP_073587.1 , 64761
Reactivity	Human
Predicted	Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	79064

PARP12 Antibody (internal region) - Additional Information

Gene ID 64761

Other Names

Poly [ADP-ribose] polymerase 12, PARP-12, 2.4.2.30, ADP-ribosyltransferase diphtheria toxin-like 12, ARTD12, Zinc finger CCCH domain-containing protein 1, PARP12, ZC3HDC1

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

PARP12 Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

PARP12 Antibody (internal region) - Protein Information

Name PARP12 {ECO:0000303|PubMed:20106667, ECO:0000312|HGNC:HGNC:21919}

Function

Mono-ADP-ribosyltransferase that mediates mono-ADP- ribosylation of target proteins.

Cellular Location

Nucleus.

PARP12 Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PARP12 Antibody (internal region) - Images**PARP12 Antibody (internal region) - References**

Identification and characterization of human TIPARP gene within the CCNL amplicon at human chromosome 3q25.31. Katoh M, Katoh M. Int J Oncol. 2003 Aug;23(2):541-7. PMID: 12851707