

Atoh7 (zebrafish) Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF2611a

Specification

Atoh7 (zebrafish) Antibody (internal region) - Product Information

Application	E
Primary Accession	O8AW52.1
Other Accession	NP_571707.1
Predicted	Zebrafish
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG

Atoh7 (zebrafish) Antibody (internal region) - Additional Information

Other Names

atoh7; atonal homolog 7 ; ath5; zath5; atonal homolog 5; lak; lakritz

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Atoh7 (zebrafish) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

Atoh7 (zebrafish) Antibody (internal region) - Protein Information

Atoh7 (zebrafish) Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Atoh7 (zebrafish) Antibody (internal region) - Images

Atoh7 (zebrafish) Antibody (internal region) - References

Staggered cell-intrinsic timing of ath5 expression underlies the wave of ganglion cell neurogenesis in the zebrafish retina. Kay JN, Link BA, Baier H. Development. 2005 Jun;132(11):2573-85. Epub 2005 Apr 27. PMID: 15857917