

BAG3 Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF2638a

Specification

BAG3 Antibody (internal region) - Product Information

Application	WB
Primary Accession	O95817
Other Accession	NP_004272.2 , 9531
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	61595

BAG3 Antibody (internal region) - Additional Information

Gene ID 9531

Other Names

BAG family molecular chaperone regulator 3, BAG-3, Bcl-2-associated athanogene 3, Bcl-2-binding protein Bis, Docking protein CAIR-1, BAG3, BIS

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

BAG3 Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

BAG3 Antibody (internal region) - Protein Information

Name BAG3

Synonyms BIS

Function

Co-chaperone for HSP70 and HSC70 chaperone proteins. Acts as a nucleotide-exchange factor (NEF) promoting the release of ADP from the HSP70 and HSC70 proteins thereby triggering client/substrate protein release. Nucleotide release is mediated via its binding to the nucleotide-binding domain (NBD) of HSPA8/HSC70 where as the substrate release is mediated via its binding to the substrate-binding domain (SBD) of HSPA8/HSC70 (PubMed:27474739, PubMed:9873016). Has anti-apoptotic activity (PubMed:10597216). Plays a role in the HSF1 nucleocytoplasmic transport (PubMed:26159920).

Cellular Location

Nucleus. Cytoplasm. Note=Colocalizes with HSF1 to the nucleus upon heat stress (PubMed:26159920)

BAG3 Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BAG3 Antibody (internal region) - Images



AF2638a (0.1 µg/ml) staining of Human Pancreas lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

BAG3 Antibody (internal region) - References

BAG3 deficiency results in fulminant myopathy and early lethality. Homma S, Iwasaki M, Shelton GD, Engvall E, Reed JC, Takayama S. Am J Pathol. 2006 Sep;169(3):761-73. PMID: 16936253