

**AOF2 Antibody (internal region)**  
Peptide-affinity purified goat antibody  
Catalog # AF2770a

**Specification**

---

**AOF2 Antibody (internal region) - Product Information**

Application	WB
Primary Accession	<a href="#">O60341</a>
Other Accession	<a href="#">NP_055828.2</a> , <a href="#">23028</a>
Reactivity	Human
Predicted	Mouse, Rat, Pig, Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	92903

**AOF2 Antibody (internal region) - Additional Information**

**Gene ID** 23028

**Other Names**

Lysine-specific histone demethylase 1A, 1.-.-., BRAF35-HDAC complex protein BHC110, Flavin-containing amine oxidase domain-containing protein 2, KDM1A, AOF2, KDM1, KIAA0601, LSD1

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

AOF2 Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

**AOF2 Antibody (internal region) - Protein Information**

**Name** KDM1A ([HGNC:29079](#))

**Function**

Histone demethylase that can demethylate both 'Lys-4' (H3K4me) and 'Lys-9' (H3K9me) of histone H3, thereby acting as a coactivator or a corepressor, depending on the context (PubMed: [15620353](http://www.uniprot.org/citations/15620353), PubMed: [15811342](http://www.uniprot.org/citations/15811342), PubMed: [16079794](http://www.uniprot.org/citations/16079794), PubMed: [16079794](#))

<http://www.uniprot.org/citations/16079795> target="\_blank">16079795</a>, PubMed:<a href="http://www.uniprot.org/citations/16140033" target="\_blank">16140033</a>, PubMed:<a href="http://www.uniprot.org/citations/16223729" target="\_blank">16223729</a>, PubMed:<a href="http://www.uniprot.org/citations/27292636" target="\_blank">27292636</a>). Acts by oxidizing the substrate by FAD to generate the corresponding imine that is subsequently hydrolyzed (PubMed:<a href="http://www.uniprot.org/citations/15620353" target="\_blank">15620353</a>, PubMed:<a href="http://www.uniprot.org/citations/15811342" target="\_blank">15811342</a>, PubMed:<a href="http://www.uniprot.org/citations/16079794" target="\_blank">16079794</a>, PubMed:<a href="http://www.uniprot.org/citations/21300290" target="\_blank">21300290</a>). Acts as a corepressor by mediating demethylation of H3K4me, a specific tag for epigenetic transcriptional activation. Demethylates both mono- (H3K4me1) and di-methylated (H3K4me2) H3K4me (PubMed:<a href="http://www.uniprot.org/citations/15620353" target="\_blank">15620353</a>, PubMed:<a href="http://www.uniprot.org/citations/20389281" target="\_blank">20389281</a>, PubMed:<a href="http://www.uniprot.org/citations/21300290" target="\_blank">21300290</a>, PubMed:<a href="http://www.uniprot.org/citations/23721412" target="\_blank">23721412</a>). May play a role in the repression of neuronal genes. Alone, it is unable to demethylate H3K4me on nucleosomes and requires the presence of RCOR1/CoREST to achieve such activity (PubMed:<a href="http://www.uniprot.org/citations/16079794" target="\_blank">16079794</a>, PubMed:<a href="http://www.uniprot.org/citations/16140033" target="\_blank">16140033</a>, PubMed:<a href="http://www.uniprot.org/citations/16885027" target="\_blank">16885027</a>, PubMed:<a href="http://www.uniprot.org/citations/21300290" target="\_blank">21300290</a>, PubMed:<a href="http://www.uniprot.org/citations/23721412" target="\_blank">23721412</a>). Also acts as a coactivator of androgen receptor (AR)-dependent transcription, by being recruited to AR target genes and mediating demethylation of H3K9me, a specific tag for epigenetic transcriptional repression. The presence of PRKCB in AR-containing complexes, which mediates phosphorylation of 'Thr-6' of histone H3 (H3T6ph), a specific tag that prevents demethylation H3K4me, prevents H3K4me demethylase activity of KDM1A (PubMed:<a href="http://www.uniprot.org/citations/16079795" target="\_blank">16079795</a>). Demethylates di-methylated 'Lys- 370' of p53/TP53 which prevents interaction of p53/TP53 with TP53BP1 and represses p53/TP53-mediated transcriptional activation. Demethylates and stabilizes the DNA methylase DNMT1 (PubMed:<a href="http://www.uniprot.org/citations/29691401" target="\_blank">29691401</a>). Demethylates methylated 'Lys-42' and methylated 'Lys-117' of SOX2 (PubMed:<a href="http://www.uniprot.org/citations/29358331" target="\_blank">29358331</a>). Required for gastrulation during embryogenesis. Component of a RCOR/GFI/KDM1A/HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development (PubMed:<a href="http://www.uniprot.org/citations/16079794" target="\_blank">16079794</a>, PubMed:<a href="http://www.uniprot.org/citations/16140033" target="\_blank">16140033</a>). Facilitates epithelial-to-mesenchymal transition by acting as an effector of SNAI1-mediated transcription repression of epithelial markers E-cadherin/CDH1, CDN7 and KRT8 (PubMed:<a href="http://www.uniprot.org/citations/20562920" target="\_blank">20562920</a>, PubMed:<a href="http://www.uniprot.org/citations/27292636" target="\_blank">27292636</a>). Required for the maintenance of the silenced state of the SNAI1 target genes E-cadherin/CDH1 and CDN7 (PubMed:<a href="http://www.uniprot.org/citations/20389281" target="\_blank">20389281</a>).

### Cellular Location

Nucleus. Chromosome. Note=Associates with chromatin

### Tissue Location

Ubiquitously expressed.


## AOF2 Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **AOF2 Antibody (internal region) - Images**



#### **AOF2 Antibody (internal region) - References**

DNA oxidation as triggered by H3K9me2 demethylation drives estrogen-induced gene expression  
Perillo B, Ombra MN, Bertoni A, Cuzzo C, Sacchetti S, Sasso A, Chiariotti L, Malorni A, Abbondanza C, Avvedimento EV Science. 2008 Jan 11;319(5860):202-6 PMID: 18187655