

**E2F8 Antibody (internal region)**  
Peptide-affinity purified goat antibody  
Catalog # AF2883a

**Specification**

---

**E2F8 Antibody (internal region) - Product Information**

Application	E
Primary Accession	<a href="#">A0AVK6</a>
Other Accession	<a href="#">NP_078956.2</a> , <a href="#">79733</a> , <a href="#">108961 (mouse)</a>
Predicted	Human, Mouse
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	94166

**E2F8 Antibody (internal region) - Additional Information**

**Gene ID** 79733

**Other Names**

Transcription factor E2F8, E2F-8, E2F8

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

E2F8 Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

**E2F8 Antibody (internal region) - Protein Information**

**Name** E2F8

**Function**

Atypical E2F transcription factor that participates in various processes such as angiogenesis and polyploidization of specialized cells. Mainly acts as a transcription repressor that binds DNA independently of DP proteins and specifically recognizes the E2 recognition site 5'-TTTC[CG]CGC-3'. Directly represses transcription of classical E2F transcription factors such as E2F1: component of a feedback loop in S phase by repressing the expression of E2F1, thereby preventing p53/TP53-dependent apoptosis. Plays a key role in polyploidization of cells in placenta and liver by regulating the endocycle, probably by repressing genes promoting cytokinesis and antagonizing action of classical E2F proteins (E2F1, E2F2 and/or E2F3). Required for placental

development by promoting polyploidization of trophoblast giant cells. Acts as a promoter of sprouting angiogenesis, possibly by acting as a transcription activator: associates with HIF1A, recognizes and binds the VEGFA promoter, which is different from canonical E2 recognition site, and activates expression of the VEGFA gene.

**Cellular Location**

Nucleus.

**E2F8 Antibody (internal region) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**E2F8 Antibody (internal region) - Images****E2F8 Antibody (internal region) - References**

Synergistic function of E2F7 and E2F8 is essential for cell survival and embryonic development. Li J, Ran C, Li E, Gordon F, Comstock G, Siddiqui H, Cleghorn W, Chen HZ, Kornacker K, Liu CG, Pandit SK, Khanizadeh M, Weinstein M, Leone G, de Bruin A. Dev. Cell 2008 Jan 14 (1): 62-75. PMID: 18194653