

**ATG4C Antibody (C- terminus)**  
Peptide-affinity purified goat antibody  
Catalog # AF2935a**Specification****ATG4C Antibody (C- terminus) - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">O96DT6</a>
Other Accession	<a href="#">NP_116241.2</a> , <a href="#">NP_835739.1</a> , <a href="#">84938</a> , <a href="#">242557</a> (mouse), <a href="#">313391</a> (rat)
Reactivity	<b>Human</b>
Predicted	<b>Dog</b>
Host	<b>Goat</b>
Clonality	<b>Polyclonal</b>
Concentration	<b>0.5 mg/ml</b>
Isotype	<b>IgG</b>
Calculated MW	<b>52497</b>

**ATG4C Antibody (C- terminus) - Additional Information****Gene ID** 84938**Other Names**

Cysteine protease ATG4C, 3.4.22.-, AUT-like 3 cysteine endopeptidase, Autophagin-3, Autophagy-related cysteine endopeptidase 3, Autophagy-related protein 4 homolog C, ATG4C, APG4C, AUTL1, AUTL3

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ATG4C Antibody (C- terminus) is for research use only and not for use in diagnostic or therapeutic procedures.

**ATG4C Antibody (C- terminus) - Protein Information****Name** ATG4C {ECO:0000303|PubMed:21177865, ECO:0000312|HGNC:HGNC:16040}**Function**

Cysteine protease that plays a key role in autophagy by mediating both proteolytic activation and delipidation of ATG8 family proteins (PubMed:&lt;a href="http://www.uniprot.org/citations/21177865" target="\_blank"&gt;21177865&lt;/a&gt;, PubMed:&lt;a href="http://www.uniprot.org/citations/29458288" target="\_blank"&gt;29458288&lt;/a&gt;, PubMed:&lt;a href="http://www.uniprot.org/citations/29458288" target="\_blank"&gt;29458288&lt;/a&gt;, PubMed:&lt;a href="http://www.uniprot.org/citations/29458288" target="\_blank"&gt;29458288&lt;/a&gt;)

<http://www.uniprot.org/citations/30661429> target="\_blank">30661429</a>). The protease activity is required for proteolytic activation of ATG8 family proteins: cleaves the C-terminal amino acid of ATG8 proteins MAP1LC3 and GABARAPL2, to reveal a C-terminal glycine (PubMed:<a href="http://www.uniprot.org/citations/21177865" target="\_blank">21177865</a>). Exposure of the glycine at the C-terminus is essential for ATG8 proteins conjugation to phosphatidylethanolamine (PE) and insertion to membranes, which is necessary for autophagy (By similarity). In addition to the protease activity, also mediates delipidation of ATG8 family proteins (PubMed:<a href="http://www.uniprot.org/citations/29458288" target="\_blank">29458288</a>, PubMed:<a href="http://www.uniprot.org/citations/33909989" target="\_blank">33909989</a>). Catalyzes delipidation of PE-conjugated forms of ATG8 proteins during macroautophagy (PubMed:<a href="http://www.uniprot.org/citations/29458288" target="\_blank">29458288</a>, PubMed:<a href="http://www.uniprot.org/citations/33909989" target="\_blank">33909989</a>). Compared to ATG4B, the major protein for proteolytic activation of ATG8 proteins, shows weaker ability to cleave the C-terminal amino acid of ATG8 proteins, while it displays stronger delipidation activity (PubMed:<a href="http://www.uniprot.org/citations/29458288" target="\_blank">29458288</a>). In contrast to other members of the family, weakly or not involved in phagophore growth during mitophagy (PubMed:<a href="http://www.uniprot.org/citations/33773106" target="\_blank">33773106</a>).

#### Cellular Location


Cytoplasm {ECO:0000250|UniProtKB:Q8BGE6}.

#### ATG4C Antibody (C- terminus) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### ATG4C Antibody (C- terminus) - Images



AF2935a (2 µg/ml) staining of Human Skeletal Muscle lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

#### **ATG4C Antibody (C- terminus) - Background**

Reported variants represent identical protein: NP\_116241.2, NP\_835739.1

#### **ATG4C Antibody (C- terminus) - References**

Six new loci associated with blood low-density lipoprotein cholesterol, high-density lipoprotein cholesterol or triglycerides in humans. Kathiresan S, Melander O, Guiducci C, Surti A, Burt NP, Rieder MJ, Cooper GM, Roos C, Voight BF, Havulinna AS, Wahlstrand B, Hedner T, Corella D, Tai ES, Ordovas JM, Berglund G, Vartiainen E, Jousilahti P, Hedblad B, Taskinen MR, Newton-Cheh C, Salomaa V Nature genetics 2008 Feb 40 (2): 189-97. PMID: 18193044