

**EDG8 / SPPR1 Antibody (internal region, near C-Term)**  
Peptide-affinity purified goat antibody  
Catalog # AF2990a

**Specification**

---

**EDG8 / SPPR1 Antibody (internal region, near C-Term) - Product Information**

Application	WB
Primary Accession	<a href="#">O9H228</a>
Other Accession	<a href="#">NP_110387.1</a> , <a href="#">53637</a>
Reactivity	Human
Predicted	Pig
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	41775

**EDG8 / SPPR1 Antibody (internal region, near C-Term) - Additional Information**

Gene ID 53637

**Other Names**

Sphingosine 1-phosphate receptor 5, S1P receptor 5, S1P5, Endothelial differentiation G-protein-coupled receptor 8, Sphingosine 1-phosphate receptor Edg-8, S1P receptor Edg-8, S1PR5, EDG8

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

EDG8 / SPPR1 Antibody (internal region, near C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

**EDG8 / SPPR1 Antibody (internal region, near C-Term) - Protein Information**

Name S1PR5

Synonyms EDG8

**Function**

Receptor for the lysosphingolipid sphingosine 1-phosphate (S1P). S1P is a bioactive lysophospholipid that elicits diverse physiological effect on most types of cells and tissues. Is coupled to both the G(i/0)alpha and G(12) subclass of heteromeric G-proteins (By similarity). May

play a regulatory role in the transformation of radial glial cells into astrocytes and may affect proliferative activity of these cells.

#### Cellular Location

Cell membrane; Multi-pass membrane protein.

#### Tissue Location


Widely expressed in the brain, most prominently in the corpus callosum, which is predominantly white matter. Detected in spleen, peripheral blood leukocytes, placenta, lung, aorta and fetal spleen. Low-level signal detected in many tissue extracts Overexpressed in leukemic large granular lymphocytes. Isoform 1 is predominantly expressed in peripheral tissues. Isoform 2 is expressed in brain, spleen and peripheral blood leukocytes

### EDG8 / SPPR1 Antibody (internal region, near C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### EDG8 / SPPR1 Antibody (internal region, near C-Term) - Images



AF2990a (1  $\mu$ g/ml) staining of Human Frontal Cortex (A) and Amygdala (B) lysates (35  $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

### EDG8 / SPPR1 Antibody (internal region, near C-Term) - References

FTY720 modulates human oligodendrocyte progenitor process extension and survival. Miron VE, Jung CG, Kim HJ, Kennedy TE, Soliven B, Antel JP. Annals of neurology 2008 Jan 63 (1): 61-71. PMID: 17918267