

**CREB3L2 / BBF2H7 Antibody (internal region)**  
**Peptide-affinity purified goat antibody**  
**Catalog # AF3078a****Specification**

---

**CREB3L2 / BBF2H7 Antibody (internal region) - Product Information**

Application	WB
Primary Accession	<a href="#">Q70SY1</a>
Other Accession	<a href="#">NP_919047.2</a> , <a href="#">64764</a> , <a href="#">208647 (mouse)</a> , <a href="#">362339 (rat)</a>
Reactivity	Mouse, Rat
Predicted	Human, Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	57415

**CREB3L2 / BBF2H7 Antibody (internal region) - Additional Information****Gene ID** 64764**Other Names**

Cyclic AMP-responsive element-binding protein 3-like protein 2, cAMP-responsive element-binding protein 3-like protein 2, BBF2 human homolog on chromosome 7, Processed cyclic AMP-responsive element-binding protein 3-like protein 2, CREB3L2, BBF2H7

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CREB3L2 / BBF2H7 Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

**CREB3L2 / BBF2H7 Antibody (internal region) - Protein Information****Name** CREB3L2**Synonyms** BBF2H7**Function**

Transcription factor involved in unfolded protein response (UPR). In the absence of endoplasmic reticulum (ER) stress, inserted into ER membranes, with N-terminal DNA-binding and transcription

activation domains oriented toward the cytosolic face of the membrane. In response to ER stress, transported to the Golgi, where it is cleaved in a site-specific manner by resident proteases S1P/MBTPS1 and S2P/MBTPS2. The released N-terminal cytosolic domain is translocated to the nucleus to effect transcription of specific target genes. Plays a critical role in chondrogenesis by activating the transcription of SEC23A, which promotes the transport and secretion of cartilage matrix proteins, and possibly that of ER biogenesis-related genes (By similarity). In a neuroblastoma cell line, protects cells from ER stress-induced death (PubMed:<a href="http://www.uniprot.org/citations/17178827" target="\_blank">17178827</a>). In vitro activates transcription of target genes via direct binding to the CRE site (PubMed:<a href="http://www.uniprot.org/citations/17178827" target="\_blank">17178827</a>).

#### Cellular Location

Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q8BH52}; Single-pass type II membrane protein Note=ER membrane resident protein. Upon ER stress, translocated to the Golgi apparatus where it is cleaved. The cytosolic N-terminal fragment (processed cyclic AMP-responsive element-binding protein 3-like protein 1) is transported into the nucleus. {ECO:0000250|UniProtKB:Q8BH52}

#### Tissue Location


Widely expressed with highest levels in placenta, lung, spleen and intestine, and lowest levels in heart, brain, skeletal muscle, thymus, colon and leukocytes. In fetal tissues, the weakest expression is detected in brain and heart

### CREB3L2 / BBF2H7 Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### CREB3L2 / BBF2H7 Antibody (internal region) - Images



AF3078a (1 µg/ml) staining of Rat Testis lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

**CREB3L2 / BBF2H7 Antibody (internal region) - References**

Characterization of the human CREB3L2 gene promoter. Panagopoulos I, Mertens F, Oncology reports 2009 Mar 21 (3): 615-24. PMID: 19212619