

BICC1 Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF3097a

Specification

BICC1 Antibody (internal region) - Product Information

Application	E
Primary Accession	O9H694
Other Accession	NP_001073981.1 , 80114 , 83675 (mouse) , 361832 (rat)
Predicted Host	Human, Mouse, Rat, Dog
Clonality	Goat
Concentration	Polyclonal
Isotype	0.5 mg/ml
Calculated MW	IgG
	104844

BICC1 Antibody (internal region) - Additional Information

Gene ID 80114

Other Names

Protein bicaudal C homolog 1, Bic-C, BICC1

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

BICC1 Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

BICC1 Antibody (internal region) - Protein Information

Name BICC1

Function

Putative RNA-binding protein. Acts as a negative regulator of Wnt signaling. May be involved in regulating gene expression during embryonic development.

Cellular Location

Cytoplasm.

BICC1 Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BICC1 Antibody (internal region) - Images**BICC1 Antibody (internal region) - References**

Association analysis of genes involved in cholesterol metabolism located within the linkage region on chromosome 10 and Alzheimer's disease. Riemenschneider M, Mahmoodzadeh S, Eisele T, Klopp N, Schwarz S, Wagenpfeil S, Diehl J, Mueller U, Foerstl H, Illig T, Kurz A, Neurobiology of aging 25 (10): 1305-8. PMID: 15465627