

**BICC1 (aa223-234) Antibody (internal region)**  
Peptide-affinity purified goat antibody  
Catalog # AF3098a**Specification**

---

**BICC1 (aa223-234) Antibody (internal region) - Product Information**

Application	WB
Primary Accession	<a href="#">O9H694</a>
Other Accession	<a href="#">NP_001073981.1</a> , <a href="#">80114</a> , <a href="#">361832 (rat)</a>
Reactivity	Human
Predicted	Rat, Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	104844

**BICC1 (aa223-234) Antibody (internal region) - Additional Information**

**Gene ID** 80114

**Other Names**

Protein bicaudal C homolog 1, Bic-C, BICC1

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

BICC1 (aa223-234) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

**BICC1 (aa223-234) Antibody (internal region) - Protein Information**

**Name** BICC1

**Function**

Putative RNA-binding protein. Acts as a negative regulator of Wnt signaling. May be involved in regulating gene expression during embryonic development.

**Cellular Location**


Cytoplasm.

## BICC1 (aa223-234) Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## BICC1 (aa223-234) Antibody (internal region) - Images



HEK293 overexpressing Human BICC1 with N-terminal FLAG, probed with rabbit anti-BICC1 in Western Blot after IP using either AF3098a or anti-FLAG antibody in the presence or absence of SDS.

## BICC1 (aa223-234) Antibody (internal region) - References

Association analysis of genes involved in cholesterol metabolism located within the linkage region on chromosome 10 and Alzheimer's disease. Riemenschneider M, Mahmoodzadeh S, Eisele T, Klopp N, Schwarz S, Wagenpfeil S, Diehl J, Mueller U, Foerstl H, Illig T, Kurz A, Neurobiology of aging 25 (10): 1305-8. PMID: 15465627