

BCL2A1 Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF3340a

Specification

BCL2A1 Antibody (internal region) - Product Information

Application	WB
Primary Accession	Q16548
Other Accession	NP_004040.1 , NP_001108207.1 , 597
Reactivity	Human
Predicted	Pig, Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	20132

BCL2A1 Antibody (internal region) - Additional Information

Gene ID 597

Other Names

Bcl-2-related protein A1, Bcl-2-like protein 5, Bcl2-L-5, Hemopoietic-specific early response protein, Protein BFL-1, Protein GRS, BCL2A1, BCL2L5, BFL1, GRS, HBPA1

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

BCL2A1 Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

BCL2A1 Antibody (internal region) - Protein Information

Name BCL2A1

Synonyms BCL2L5, BFL1, GRS, HBPA1

Function

Retards apoptosis induced by IL-3 deprivation. May function in the response of hemopoietic cells to external signals and in maintaining endothelial survival during infection (By similarity). Can inhibit apoptosis induced by serum starvation in the mammary epithelial cell line HC11 (By similarity).

Cellular Location

Cytoplasm.

Tissue Location

Seems to be restricted to the hematopoietic compartment. Expressed in peripheral blood, spleen, and bone marrow, at moderate levels in lung, small intestine and testis, at a minimal levels in other tissues. Also found in vascular smooth muscle cells and hematopoietic malignancies

BCL2A1 Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BCL2A1 Antibody (internal region) - Images

AF3304a (0.2 µg/ml) staining of Human Bone Marrow lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

BCL2A1 Antibody (internal region) - Background

This antibody is expected to recognize both reported isoforms (NP_004040.1; NP_001108207.1).

BCL2A1 Antibody (internal region) - References

Association between genetic variants in VEGF, ERCC3 and occupational benzene haematotoxicity. Hosgood HD, Zhang L, Shen M, Berndt SI, Vermeulen R, Li G, Yin S, Yeager M, Yuenger J, Rothman N, Chanock S, Smith M, Lan Q, Occupational and environmental medicine 2009 Dec 66 (12): 848-53. PMID: 19773279