

Alk (mouse) Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF3598a**Specification****Alk (mouse) Antibody (internal region) - Product Information**

Application	WB
Primary Accession	O9UM73
Other Accession	NP_031465.2 , 238 , 11682 (mouse) , 266802 (rat)
Reactivity	Rat
Predicted	Mouse
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	176442

Alk (mouse) Antibody (internal region) - Additional Information**Gene ID** 238**Other Names**

ALK tyrosine kinase receptor, 2.7.10.1, Anaplastic lymphoma kinase, CD246, ALK

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Alk (mouse) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

Alk (mouse) Antibody (internal region) - Protein Information**Name** ALK {ECO:0000303|PubMed:9174053, ECO:0000312|HGNC:HGNC:427}**Function**

Neuronal receptor tyrosine kinase that is essentially and transiently expressed in specific regions of the central and peripheral nervous systems and plays an important role in the genesis and differentiation of the nervous system (PubMed:11121404, PubMed:11387242, PubMed:16317043, PubMed:16317043, PubMed:16317043)

[17274988](http://www.uniprot.org/citations/17274988), PubMed: [30061385](http://www.uniprot.org/citations/30061385), PubMed: [34646012](http://www.uniprot.org/citations/34646012), PubMed: [34819673](http://www.uniprot.org/citations/34819673)). Also acts as a key thinness protein involved in the resistance to weight gain: in hypothalamic neurons, controls energy expenditure acting as a negative regulator of white adipose tissue lipolysis and sympathetic tone to fine-tune energy homeostasis (By similarity). Following activation by ALKAL2 ligand at the cell surface, transduces an extracellular signal into an intracellular response (PubMed: [30061385](http://www.uniprot.org/citations/30061385), PubMed: [33411331](http://www.uniprot.org/citations/33411331), PubMed: [34646012](http://www.uniprot.org/citations/34646012), PubMed: [34819673](http://www.uniprot.org/citations/34819673)). In contrast, ALKAL1 is not a potent physiological ligand for ALK (PubMed: [34646012](http://www.uniprot.org/citations/34646012)). Ligand-binding to the extracellular domain induces tyrosine kinase activation, leading to activation of the mitogen-activated protein kinase (MAPK) pathway (PubMed: [34819673](http://www.uniprot.org/citations/34819673)). Phosphorylates almost exclusively at the first tyrosine of the Y-x-x-x-Y-Y motif (PubMed: [15226403](http://www.uniprot.org/citations/15226403), PubMed: [16878150](http://www.uniprot.org/citations/16878150)). Induces tyrosine phosphorylation of CBL, FRS2, IRS1 and SHC1, as well as of the MAP kinases MAPK1/ERK2 and MAPK3/ERK1 (PubMed: [15226403](http://www.uniprot.org/citations/15226403), PubMed: [16878150](http://www.uniprot.org/citations/16878150)). ALK activation may also be regulated by pleiotrophin (PTN) and midkine (MDK) (PubMed: [11278720](http://www.uniprot.org/citations/11278720), PubMed: [11809760](http://www.uniprot.org/citations/11809760), PubMed: [12107166](http://www.uniprot.org/citations/12107166), PubMed: [12122009](http://www.uniprot.org/citations/12122009)). PTN-binding induces MAPK pathway activation, which is important for the anti-apoptotic signaling of PTN and regulation of cell proliferation (PubMed: [11278720](http://www.uniprot.org/citations/11278720), PubMed: [11809760](http://www.uniprot.org/citations/11809760), PubMed: [12107166](http://www.uniprot.org/citations/12107166)). MDK-binding induces phosphorylation of the ALK target insulin receptor substrate (IRS1), activates mitogen-activated protein kinases (MAPKs) and PI3-kinase, resulting also in cell proliferation induction (PubMed: [12122009](http://www.uniprot.org/citations/12122009)). Drives NF-kappa-B activation, probably through IRS1 and the activation of the AKT serine/threonine kinase (PubMed: [15226403](http://www.uniprot.org/citations/15226403), PubMed: [16878150](http://www.uniprot.org/citations/16878150)). Recruitment of IRS1 to activated ALK and the activation of NF-kappa-B are essential for the autocrine growth and survival signaling of MDK (PubMed: [15226403](http://www.uniprot.org/citations/15226403), PubMed: [16878150](http://www.uniprot.org/citations/16878150)).

Cellular Location

Cell membrane; Single-pass type I membrane protein Note=Membrane attachment is essential for promotion of neuron-like differentiation and cell proliferation arrest through specific activation of the MAP kinase pathway.

Tissue Location


Expressed in brain and CNS. Also expressed in the small intestine and testis, but not in normal lymphoid cells

Alk (mouse) Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Alk (mouse) Antibody (internal region) - Images



Alk (mouse) Antibody (internal region) - References

The neuroblastoma-associated F1174L ALK mutation causes resistance to an ALK kinase inhibitor in ALK-translocated cancers. Sasaki T, Okuda K, Zheng W, Butrynski J, Capelletti M, Wang L, Gray NS, Wilner K, Christensen JG, Demetri G, Shapiro GI, Rodig SJ, Eck MJ, Jänne PA. Cancer Res. 2010 Dec 15;70(24):10038-43. PMID: 21030459