

Atrial natriuretic factor Antibody (C-Term)
Peptide-affinity purified goat antibody
Catalog # AF3647a

Specification

Atrial natriuretic factor Antibody (C-Term) - Product Information

Application	IHC, WB
Primary Accession	P01160
Other Accession	NP_006163.1 , 4878 , 230899 (mouse) , 24602 (rat)
Reactivity	Human, Mouse, Rat
Predicted	Pig, Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	16396

Atrial natriuretic factor Antibody (C-Term) - Additional Information

Gene ID 4878

Other Names

Natriuretic peptides A, CDD-ANF, Cardiodilatin, CDD, Cardiodilatin-related peptide, CDP, Prepronatriodilatin, Atrial natriuretic factor, ANF, Atrial natriuretic peptide, ANP, NPPA, ANP, PND

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Atrial natriuretic factor Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

Atrial natriuretic factor Antibody (C-Term) - Protein Information

Name NPPA

Synonyms ANP, PND

Function

[Atrial natriuretic peptide]: Hormone that plays a key role in mediating cardio-renal homeostasis, and is involved in vascular remodeling and regulating energy metabolism (PubMed:15741263, PubMed:16875975, PubMed:18835931, PubMed:21672517, PubMed:22307324, PubMed:2532366, PubMed:2825692, PubMed:7595132, PubMed:7720651, PubMed:8087923, PubMed:8653797). Acts by specifically binding and stimulating NPR1 to produce cGMP, which in turn activates effector proteins, such as PRKG1, that drive various biological responses (PubMed:1660465, PubMed:1672777, PubMed:21098034, PubMed:2162527, PubMed:22307324, PubMed:25401746, PubMed:2825692, PubMed:7720651, PubMed:8384600, PubMed:9893117). Regulates vasodilation, natriuresis, diuresis and aldosterone synthesis and is therefore essential for regulating blood pressure, controlling the extracellular fluid volume and maintaining the fluid-electrolyte balance (PubMed:2532366, PubMed:2825692, PubMed:7595132, PubMed:7720651, PubMed:8087923, PubMed:8653797). Also involved in inhibiting cardiac remodeling and cardiac hypertrophy by inducing cardiomyocyte apoptosis and attenuating the growth of cardiomyocytes and fibroblasts (PubMed:16875975). Plays a role in female pregnancy by promoting trophoblast invasion and spiral artery remodeling in uterus, and thus prevents pregnancy-induced hypertension (By similarity). In adipose tissue, acts in various cGMP- and PKG-dependent pathways to regulate lipid metabolism and energy homeostasis (PubMed:15741263, PubMed:18835931, PubMed:21672517, PubMed:22307324). This includes up-regulating lipid metabolism and mitochondrial oxygen utilization by activating the AMP-activated protein kinase (AMPK), and increasing energy expenditure by acting via MAPK11 to promote the UCP1-dependent thermogenesis of brown adipose tissue (PubMed:15741263, PubMed:18835931, PubMed:21672517, PubMed:22307324). Binds the clearance receptor NPR3 which removes the hormone from circulation (PubMed:1672777).

Cellular Location

[Long-acting natriuretic peptide]: Secreted. Note=Detected in blood. [Kaliuretic peptide]: Secreted. Note=Detected in blood [Atrial natriuretic peptide]: Secreted. Perikaryon. Cell projection. Note=Detected in blood (PubMed:15741263, PubMed:18835931, PubMed:2532366, PubMed:7955907, PubMed:7984506, PubMed:8351194, PubMed:8653797, PubMed:8779891). Detected in urine in one study (PubMed:8351194). However, in another study, was not detected in

urine (PubMed:7984506). Detected in cytoplasmic bodies and neuronal processes of pyramidal neurons (layers II-VI) (PubMed:30534047) Increased secretion in response to the vasopressin AVP (By similarity) Likely to be secreted in response to an increase in atrial pressure or atrial stretch (PubMed:2532366). In kidney cells, secretion increases in response to activated guanylyl cyclases and increased intracellular cAMP levels (PubMed:9893117). Plasma levels increase 15 minutes after a high-salt meal, and decrease back to normal plasma levels 1 hr later (PubMed:8779891). {ECO:0000250|UniProtKB:P01161, ECO:0000269|PubMed:15741263, ECO:0000269|PubMed:18835931, ECO:0000269|PubMed:2532366, ECO:0000269|PubMed:30534047, ECO:0000269|PubMed:7955907, ECO:0000269|PubMed:7984506, ECO:0000269|PubMed:8351194, ECO:0000269|PubMed:8653797, ECO:0000269|PubMed:8779891, ECO:0000269|PubMed:9893117}

Tissue Location


[Urodilatin]: Detected in the kidney distal tubular cells (at protein level) (PubMed:8384600, PubMed:9794555). Present in urine (at protein level) (PubMed:2972874, PubMed:8351194, PubMed:8779891, PubMed:9794555).

Atrial natriuretic factor Antibody (C-Term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Atrial natriuretic factor Antibody (C-Term) - Images



AF3647a (5 µg/ml) staining of paraffin embedded Human Heart. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



AF3647a (0.01 $\mu\text{g/ml}$) staining of Mouse Heart lysate (35 μg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

Atrial natriuretic factor Antibody (C-Term) - Background

The immunizing peptide represents aa137-150 of the precursor NPPA.

Atrial natriuretic factor Antibody (C-Term) - References

Decompensated heart failure is associated with reduced corin levels and decreased cleavage of pro-atrial natriuretic peptide. Ibebuogu UN, Gladysheva IP, Houg AK, Reed GL. *Circ Heart Fail.* 2011 Mar 1;4(2):114-20. PMID: 21216831