

**ARNT (aa558-570) Antibody (internal region)**  
Peptide-affinity purified goat antibody  
Catalog # AF3670a

### Specification

---

#### ARNT (aa558-570) Antibody (internal region) - Product Information

Application	WB
Primary Accession	<a href="#">P27540</a>
Other Accession	<a href="#">NP_001659.1</a> , <a href="#">NP_848514.1</a> , <a href="#">NP_001184254.1</a> , <a href="#">405</a>
Reactivity	Human
Predicted	Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	86636

#### ARNT (aa558-570) Antibody (internal region) - Additional Information

Gene ID 405

#### Other Names

Aryl hydrocarbon receptor nuclear translocator, ARNT protein, Class E basic helix-loop-helix protein 2, bHLHe2, Dioxin receptor, nuclear translocator, Hypoxia-inducible factor 1-beta, HIF-1-beta, HIF1-beta, ARNT, BHLHE2

#### Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

ARNT (aa558-570) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

#### ARNT (aa558-570) Antibody (internal region) - Protein Information

Name ARNT ([HGNC:700](#))

Synonyms BHLHE2

#### Function

Required for activity of the AHR. Upon ligand binding, AHR translocates into the nucleus, where it heterodimerizes with ARNT and induces transcription by binding to xenobiotic response elements

(XRE). Not required for the ligand-binding subunit to translocate from the cytosol to the nucleus after ligand binding (PubMed:<a href="http://www.uniprot.org/citations/34521881" target="\_blank">34521881</a>). The complex initiates transcription of genes involved in the regulation of a variety of biological processes, including angiogenesis, hematopoiesis, drug and lipid metabolism, cell motility and immune modulation (Probable). The heterodimer binds to core DNA sequence 5'-TACGTG-3' within the hypoxia response element (HRE) of target gene promoters and functions as a transcriptional regulator of the adaptive response to hypoxia (By similarity). The heterodimer ARNT:AHR binds to core DNA sequence 5'-TGCGTG-3' within the dioxin response element (DRE) of target gene promoters and activates their transcription (PubMed:<a href="http://www.uniprot.org/citations/28396409" target="\_blank">28396409</a>).

#### Cellular Location

Nucleus.

#### ARNT (aa558-570) Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### ARNT (aa558-570) Antibody (internal region) - Images



AF3670a (1 µg/ml) staining of HeLa nuclear lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

#### ARNT (aa558-570) Antibody (internal region) - Background

This antibody is expected to recognize all reported isoforms (NP\_001659.1; NP\_848514.1; NP\_001184254.1).

**ARNT (aa558-570) Antibody (internal region) - References**

ETV6-ARNT fusion in a patient with childhood T lymphoblastic leukemia. Otsubo K, Kanegane H, Eguchi M, Eguchi-Ishimae M, Tamura K, Nomura K, Abe A, Ishii E, Miyawaki T. *Cancer Genet Cytogenet.* 2010 Oct 1;202(1):22-6. PMID: 20804916