

CP110 Antibody (N-Term)
Peptide-affinity purified goat antibody
Catalog # AF3694a

Specification

CP110 Antibody (N-Term) - Product Information

Application	WB
Primary Accession	O43303
Other Accession	NP_001185951.1 , NP_055526.3 , 9738
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	113424

CP110 Antibody (N-Term) - Additional Information

Gene ID 9738

Other Names

Centriolar coiled-coil protein of 110 kDa, Centrosomal protein of 110 kDa, CP110, Cep110, CCP110, CEP110, CP110, KIAA0419

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CP110 Antibody (N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

CP110 Antibody (N-Term) - Protein Information

Name CCP110

Synonyms CEP110, CP110, KIAA0419

Function

Necessary for centrosome duplication at different stages of procentriole formation. Acts as a key negative regulator of ciliogenesis in collaboration with CEP97 by capping the mother centriole thereby preventing cilia formation (PubMed: [17681131](http://www.uniprot.org/citations/17681131), PubMed: [17719545](http://www.uniprot.org/citations/17719545), PubMed: [23486064](http://www.uniprot.org/citations/23486064))

target="_blank">23486064, PubMed:30375385, PubMed:35301795). Also involved in promoting ciliogenesis. May play a role in the assembly of the mother centriole subdistal appendages (SDA) thereby effecting the fusion of recycling endosomes to basal bodies during cilia formation (By similarity). Required for correct spindle formation and has a role in regulating cytokinesis and genome stability via cooperation with CALM1 and CETN2 (PubMed:16760425).

Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, cilium basal body {ECO:0000250|UniProtKB:Q7TSH4} Note=Recruited early and then associates with the growing distal tips Recruited to the mother centriole by KIF24 (PubMed:21620453). Removed from centrioles by TTBK2, leading to initiation of ciliogenesis and localizes only to the daughter centriole in ciliated cells. In cytotoxic T lymphocytes remains associated with the mother centriole during docking of the centrosome at the immunological synapse upon target contact (By similarity). Recruited at the distal end of the mother centriole by MPHOSPH9 (PubMed:30375385) {ECO:0000250|UniProtKB:Q7TSH4, ECO:0000269|PubMed:21620453, ECO:0000269|PubMed:30375385}

Tissue Location

Highly expressed in testis. Detected at intermediate levels in spleen, thymus, prostate, small intestine, colon and peripheral blood leukocytes.


CP110 Antibody (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CP110 Antibody (N-Term) - Images





AF3694a (1 μ g/ml) staining of Human Testis lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

CP110 Antibody (N-Term) - Background

This antibody is expected to recognize both reported isoforms (NP_001185951.1; NP_055526.3).

CP110 Antibody (N-Term) - References

SCF(Cyclin F) controls centrosome homeostasis and mitotic fidelity through CP110 degradation.
D'Angiolella V, Donato V, Vijayakumar S, Saraf A, Florens L, Washburn MP, Dynlacht B, Pagano M.
Nature. 2010 Jul 1;466(7302):138-42. PMID: 20596027