

ROS1 Antibody (C-Term)
Peptide-affinity purified goat antibody
Catalog # AF3777a

Specification

ROS1 Antibody (C-Term) - Product Information

Application	E
Primary Accession	P08922
Other Accession	NP_002935.2 , 6098
Predicted	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	263915

ROS1 Antibody (C-Term) - Additional Information

Gene ID 6098

Other Names

Proto-oncogene tyrosine-protein kinase ROS, 2.7.10.1, Proto-oncogene c-Ros, Proto-oncogene c-Ros-1, Receptor tyrosine kinase c-ros oncogene 1, c-Ros receptor tyrosine kinase, ROS1, MCF3, ROS

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ROS1 Antibody (C-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

ROS1 Antibody (C-Term) - Protein Information

Name ROS1

Synonyms MCF3, ROS

Function

Receptor tyrosine kinase (RTK) that plays a role in epithelial cell differentiation and regionalization of the proximal epididymal epithelium. NELL2 is an endogenous ligand for ROS1. Upon endogenous stimulation by NELL2, ROS1 activates the intracellular signaling pathway and triggers epididymal epithelial differentiation and subsequent sperm maturation (By similarity). May

activate several downstream signaling pathways related to cell differentiation, proliferation, growth and survival including the PI3 kinase-mTOR signaling pathway. Mediates the phosphorylation of PTPN11, an activator of this pathway. May also phosphorylate and activate the transcription factor STAT3 to control anchorage-independent cell growth. Mediates the phosphorylation and the activation of VAV3, a guanine nucleotide exchange factor regulating cell morphology. May activate other downstream signaling proteins including AKT1, MAPK1, MAPK3, IRS1 and PLCG2.

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location

Expressed in brain. Expression is increased in primary gliomas.

ROS1 Antibody (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ROS1 Antibody (C-Term) - Images**ROS1 Antibody (C-Term) - References**

Identification of four gene variants associated with myocardial infarction. Shiffman D, Ellis SG, Rowland CM, Malloy MJ, Luke MM, Iakoubova OA, Pullinger CR, Cassano J, Aouizerat BE, Fenwick RG, Reitz RE, Catanese JJ, Leong DU, Zellner C, Sninsky JJ, Topol EJ, Devlin JJ, Kane JP. Am J Hum Genet. 2005 Oct;77(4):596-605. PMID: 16175505