

LSM2 (aa33-46) Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF3784a**Specification**

LSM2 (aa33-46) Antibody (internal region) - Product Information

Application	WB
Primary Accession	O9Y333
Other Accession	NP_067000.1 , 57819 , 27756 (mouse) , 684148 (rat)
Reactivity	Human, Mouse
Predicted	Rat, Pig, Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	10835

LSM2 (aa33-46) Antibody (internal region) - Additional Information

Gene ID 57819

Other Names

U6 snRNA-associated Sm-like protein LSm2, Protein G7b, Small nuclear ribonuclear protein D homolog, snRNP core Sm-like protein Sm-x5, LSM2, C6orf28, G7B

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

LSM2 (aa33-46) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

LSM2 (aa33-46) Antibody (internal region) - Protein Information

Name LSM2

Synonyms C6orf28, G7B

Function

Plays a role in pre-mRNA splicing as component of the U4/U6- U5 tri-snRNP complex that is involved in spliceosome assembly, and as component of the precatalytic spliceosome (spliceosome B complex) (PubMed:

target="_blank">28781166). The heptameric LSM2-8 complex binds specifically to the 3'-terminal U-tract of U6 snRNA (PubMed:10523320).

Cellular Location

Nucleus

LSM2 (aa33-46) Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

LSM2 (aa33-46) Antibody (internal region) - Images



AF3784a (1 µg/ml) staining of K562 lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

LSM2 (aa33-46) Antibody (internal region) - References

The human LSm1-7 proteins colocalize with the mRNA-degrading enzymes Dcp1/2 and Xrnl in distinct cytoplasmic foci. Ingelfinger D, Arndt-Jovin DJ, Lührmann R, Achsel T. RNA. 2002 Dec;8(12):1489-501. PMID: 12515382