

**TUBB3 Antibody (internal region, near N-Term)**  
Peptide-affinity purified goat antibody  
Catalog # AF3836a

**Specification**

---

**TUBB3 Antibody (internal region, near N-Term) - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">Q13509</a>
Other Accession	<a href="#">NP_006077.2</a> , <a href="#">10381</a> , <a href="#">22152 (mouse)</a> , <a href="#">246118 (rat)</a>
Reactivity	<b>Human</b>
Predicted	<b>Mouse, Rat, Pig</b>
Host	<b>Goat</b>
Clonality	<b>Polyclonal</b>
Concentration	<b>0.5 mg/ml</b>
Isotype	<b>IgG</b>
Calculated MW	<b>50433</b>

**TUBB3 Antibody (internal region, near N-Term) - Additional Information**

**Gene ID** 10381

**Other Names**

Tubulin beta-3 chain, Tubulin beta-4 chain, Tubulin beta-III, TUBB3, TUBB4

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

TUBB3 Antibody (internal region, near N-Term) is for research use only and not for use in diagnostic or therapeutic procedures.

**TUBB3 Antibody (internal region, near N-Term) - Protein Information**

**Name** TUBB3

**Synonyms** TUBB4

**Function**

Tubulin is the major constituent of microtubules, a cylinder consisting of laterally associated linear protofilaments composed of alpha- and beta-tubulin heterodimers (PubMed: [34996871](http://www.uniprot.org/citations/34996871)). Microtubules grow by the addition of GTP-tubulin dimers to the microtubule end, where a stabilizing cap forms

(PubMed:<a href="http://www.uniprot.org/citations/34996871" target="\_blank">34996871</a>). Below the cap, tubulin dimers are in GDP-bound state, owing to GTPase activity of alpha- tubulin (PubMed:<a href="http://www.uniprot.org/citations/34996871" target="\_blank">34996871</a>). TUBB3 plays a critical role in proper axon guidance and maintenance (PubMed:<a href="http://www.uniprot.org/citations/20074521" target="\_blank">20074521</a>). Binding of NTN1/Netrin-1 to its receptor UNC5C might cause dissociation of UNC5C from polymerized TUBB3 in microtubules and thereby lead to increased microtubule dynamics and axon repulsion (PubMed:<a href="http://www.uniprot.org/citations/28483977" target="\_blank">28483977</a>). Plays a role in dorsal root ganglion axon projection towards the spinal cord (PubMed:<a href="http://www.uniprot.org/citations/28483977" target="\_blank">28483977</a>).

#### Cellular Location

Cytoplasm, cytoskeleton. Cell projection, growth cone {ECO:0000250|UniProtKB:Q9ERD7}. Cell projection, lamellipodium {ECO:0000250|UniProtKB:Q9ERD7}. Cell projection, filopodium {ECO:0000250|UniProtKB:Q9ERD7}

#### Tissue Location

Expression is primarily restricted to central and peripheral nervous system. Greatly increased expression in most cancerous tissues.

#### TUBB3 Antibody (internal region, near N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### TUBB3 Antibody (internal region, near N-Term) - Images



AF3836a (2 µg/ml) staining of HepG2 lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

**TUBB3 Antibody (internal region, near N-Term) - Background**

This antibody is expected to recognize isoform 1 (NP\_006077.2) only.

**TUBB3 Antibody (internal region, near N-Term) - References**

Class III  $\beta$ -tubulin expression in advanced-stage serous ovarian carcinoma effusions is associated with poor survival and primary chemoresistance. Hetland TE, Hellesylt E, Flørenes VA, Tropé C, Davidson B, Kærn J. Hum Pathol. 2011 Jul;42(7):1019-26. PMID: 21315408