

**Cytokeratin 20, Biotinylated**  
Peptide-affinity purified goat antibody  
Catalog # AF3856b

**Specification**

---

**Cytokeratin 20, Biotinylated - Product Information**

Application	WB
Primary Accession	<a href="#">P35900</a>
Other Accession	<a href="#">NP_061883.1</a> , <a href="#">54474</a>
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	48487

**Cytokeratin 20, Biotinylated - Additional Information**

**Gene ID** 54474

**Other Names**

Keratin, type I cytoskeletal 20, Cytokeratin-20, CK-20, Keratin-20, K20, Protein IT, KRT20

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Cytokeratin 20, Biotinylated is for research use only and not for use in diagnostic or therapeutic procedures.

**Cytokeratin 20, Biotinylated - Protein Information**

**Name** KRT20

**Function**

Plays a significant role in maintaining keratin filament organization in intestinal epithelia. When phosphorylated, plays a role in the secretion of mucin in the small intestine (By similarity).

**Cellular Location**

Cytoplasm

**Tissue Location**

Expressed predominantly in the intestinal epithelium. Expressed in luminal cells of colonic mucosa.

Also expressed in the Merkel cells of keratinized oral mucosa; specifically at the tips of some rete ridges of the gingival mucosa, in the basal layer of the palatal mucosa and in the taste buds of lingual mucosa

### Cytokeratin 20, Biotinylated - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Cytokeratin 20, Biotinylated - Images



Biotinylated EB11745 (1 $\mu$ g/ml) staining of Human Colon lysate (35 $\mu$ g protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker

### Cytokeratin 20, Biotinylated - References

SATB2 in combination with cytokeratin 20 identifies over 95% of all colorectal carcinomas. Magnusson K, de Wit M, Brennan DJ, Johnson LB, McGee SF, Lundberg E, Naicker K, Klinger R, Kampf C, Asplund A, Wester K, Gry M, Bjartell A, Gallagher WM, Rexhepaj E, Kilpinen S, Kallioniemi OP, Belt E, Goos J, Meijer G, Birgisson H, Glimelius B, Borrebaec Am J Surg Pathol. 2011 Jul;35(7):937-48. PMID: 21677534