

**IL-17B (aa77-89) Antibody (internal region)**  
Peptide-affinity purified goat antibody  
Catalog # AF3981a

**Specification**

---

**IL-17B (aa77-89) Antibody (internal region) - Product Information**

Application	WB
Primary Accession	<a href="#">O9UHF5</a>
Other Accession	<a href="#">NP_055258.1</a> , <a href="#">27190</a>
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	20437

**IL-17B (aa77-89) Antibody (internal region) - Additional Information**

**Gene ID** 27190

**Other Names**

Interleukin-17B, IL-17B, Cytokine Zcyto7, Interleukin-20, IL-20, Neuronal interleukin-17-related factor, IL17B, IL20, NIRF, ZCYTO7

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

IL-17B (aa77-89) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

**IL-17B (aa77-89) Antibody (internal region) - Protein Information**

**Name** IL17B

**Synonyms** IL20, NIRF, ZCYTO7

**Function**

Stimulates the release of tumor necrosis factor alpha and IL- 1-beta from the monocytic cell line THP-1.

**Cellular Location**

Secreted.

**Tissue Location**

Expressed in adult pancreas, small intestine, stomach, spinal cord and testis. Less pronounced expression in prostate, colon mucosal lining, and ovary

**IL-17B (aa77-89) Antibody (internal region) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**IL-17B (aa77-89) Antibody (internal region) - Images**

AF3981a (2 µg/ml) staining of Human Ileum lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

**IL-17B (aa77-89) Antibody (internal region) - References**

IL-17B and IL-17C are associated with TNF-alpha production and contribute to the exacerbation of inflammatory arthritis. Yamaguchi Y, Fujio K, Shoda H, Okamoto A, Tsuno NH, Takahashi K, Yamamoto K. J Immunol. 2007 Nov 15;179(10):7128-36. PMID: 17982105