

**IFNAR1 (aa353-364) Antibody (internal region)**  
Peptide-affinity purified goat antibody  
Catalog # AF3984a

### Specification

---

#### IFNAR1 (aa353-364) Antibody (internal region) - Product Information

Application	WB
Primary Accession	<a href="#">P17181</a>
Other Accession	<a href="#">NP_000620.2</a> , <a href="#">3454</a>
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	63525

#### IFNAR1 (aa353-364) Antibody (internal region) - Additional Information

Gene ID 3454

#### Other Names

Interferon alpha/beta receptor 1, IFN-R-1, IFN-alpha/beta receptor 1, Cytokine receptor class-II member 1, Cytokine receptor family 2 member 1, CRF2-1, Type I interferon receptor 1, IFNAR1, IFNAR

#### Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

IFNAR1 (aa353-364) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

#### IFNAR1 (aa353-364) Antibody (internal region) - Protein Information

Name IFNAR1

Synonyms IFNAR

#### Function

Together with IFNAR2, forms the heterodimeric receptor for type I interferons (including interferons alpha, beta, epsilon, omega and kappa) (PubMed:<a href="http://www.uniprot.org/citations/10049744" target="\_blank">10049744</a>, PubMed:<a href="http://www.uniprot.org/citations/14532120" target="\_blank">14532120</a>, PubMed:<a href="http://www.uniprot.org/citations/14532120" target="\_blank">14532120</a>, PubMed:<a href="http://www.uniprot.org/citations/14532120" target="\_blank">14532120</a>)

<http://www.uniprot.org/citations/15337770> target="\_blank">15337770</a>, PubMed:<a href="http://www.uniprot.org/citations/2153461" target="\_blank">2153461</a>, PubMed:<a href="http://www.uniprot.org/citations/21854986" target="\_blank">21854986</a>, PubMed:<a href="http://www.uniprot.org/citations/24075985" target="\_blank">24075985</a>, PubMed:<a href="http://www.uniprot.org/citations/31270247" target="\_blank">31270247</a>, PubMed:<a href="http://www.uniprot.org/citations/33252644" target="\_blank">33252644</a>, PubMed:<a href="http://www.uniprot.org/citations/35442418" target="\_blank">35442418</a>, PubMed:<a href="http://www.uniprot.org/citations/7813427" target="\_blank">7813427</a>). Type I interferon binding activates the JAK-STAT signaling cascade, resulting in transcriptional activation or repression of interferon-regulated genes that encode the effectors of the interferon response (PubMed:<a href="http://www.uniprot.org/citations/10049744" target="\_blank">10049744</a>, PubMed:<a href="http://www.uniprot.org/citations/21854986" target="\_blank">21854986</a>, PubMed:<a href="http://www.uniprot.org/citations/7665574" target="\_blank">7665574</a>). Mechanistically, type I interferon- binding brings the IFNAR1 and IFNAR2 subunits into close proximity with one another, driving their associated Janus kinases (JAKs) (TYK2 bound to IFNAR1 and JAK1 bound to IFNAR2) to cross-phosphorylate one another (PubMed:<a href="http://www.uniprot.org/citations/21854986" target="\_blank">21854986</a>, PubMed:<a href="http://www.uniprot.org/citations/32972995" target="\_blank">32972995</a>, PubMed:<a href="http://www.uniprot.org/citations/7665574" target="\_blank">7665574</a>, PubMed:<a href="http://www.uniprot.org/citations/7813427" target="\_blank">7813427</a>). The activated kinases phosphorylate specific tyrosine residues on the intracellular domains of IFNAR1 and IFNAR2, forming docking sites for the STAT transcription factors (PubMed:<a href="http://www.uniprot.org/citations/21854986" target="\_blank">21854986</a>, PubMed:<a href="http://www.uniprot.org/citations/32972995" target="\_blank">32972995</a>, PubMed:<a href="http://www.uniprot.org/citations/7526154" target="\_blank">7526154</a>, PubMed:<a href="http://www.uniprot.org/citations/7665574" target="\_blank">7665574</a>, PubMed:<a href="http://www.uniprot.org/citations/7813427" target="\_blank">7813427</a>). STAT proteins are then phosphorylated by the JAKs, promoting their translocation into the nucleus to regulate expression of interferon-regulated genes (PubMed:<a href="http://www.uniprot.org/citations/19561067" target="\_blank">19561067</a>, PubMed:<a href="http://www.uniprot.org/citations/21854986" target="\_blank">21854986</a>, PubMed:<a href="http://www.uniprot.org/citations/32972995" target="\_blank">32972995</a>, PubMed:<a href="http://www.uniprot.org/citations/7665574" target="\_blank">7665574</a>, PubMed:<a href="http://www.uniprot.org/citations/7813427" target="\_blank">7813427</a>, PubMed:<a href="http://www.uniprot.org/citations/9121453" target="\_blank">9121453</a>). Can also act independently of IFNAR2: form an active IFNB1 receptor by itself and activate a signaling cascade that does not involve activation of the JAK-STAT pathway (By similarity).

### Cellular Location

[Isoform 1]: Cell membrane; Single-pass type I membrane protein. Late endosome. Lysosome. Note=Interferon binding triggers internalization of the receptor from the cell membrane into endosomes and then into lysosomes.

### Tissue Location

IFN receptors are present in all tissues and even on the surface of most IFN-resistant cells. Isoform 1, isoform 2 and isoform 3 are expressed in the IFN-alpha sensitive myeloma cell line U266B1. Isoform 2 and isoform 3 are expressed in the IFN-alpha resistant myeloma cell line U266R. Isoform 1 is not expressed in IFN- alpha resistant myeloma cell line U266R.


### IFNAR1 (aa353-364) Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **IFNAR1 (aa353-364) Antibody (internal region) - Images**



#### **IFNAR1 (aa353-364) Antibody (internal region) - References**

Type I IFN-dependent T cell activation is mediated by IFN-dependent dendritic cell OX40 ligand expression and is independent of T cell IFNR expression. Kurche JS, Haluszczak C, McWilliams JA, Sanchez PJ, Kedl RM. J Immunol. 2012 Jan 15;188(2):585-93. PMID: 22156349