

cytokeratin 19 (aa285-298) Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF4070a

Specification

cytokeratin 19 (aa285-298) Antibody (internal region) - Product Information

Application	WB, IHC
Primary Accession	P08727
Other Accession	NP_002267.2 , 3880
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	44106

cytokeratin 19 (aa285-298) Antibody (internal region) - Additional Information

Gene ID 3880

Other Names

Keratin, type I cytoskeletal 19, Cytokeratin-19, CK-19, Keratin-19, K19, KRT19

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

cytokeratin 19 (aa285-298) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

cytokeratin 19 (aa285-298) Antibody (internal region) - Protein Information

Name KRT19

Function

Involved in the organization of myofibers. Together with KRT8, helps to link the contractile apparatus to dystrophin at the costameres of striated muscle.

Tissue Location

Expressed in a defined zone of basal keratinocytes in the deep outer root sheath of hair follicles. Also observed in sweat gland and mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, ectocervical epithelium (at protein level). Expressed in epidermal basal cells, in nipple epidermis and a defined region of the


hair follicle. Also seen in a subset of vascular wall cells in both the veins and artery of human umbilical cord, and in umbilical cord vascular smooth muscle. Observed in muscle fibers accumulating in the costameres of myoplasm at the sarcolemma in structures that contain dystrophin and spectrin.

cytokeratin 19 (aa285-298) Antibody (internal region) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

cytokeratin 19 (aa285-298) Antibody (internal region) - Images



E12359 (0.03 μ g/ml) staining of Human Colon cancer lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



AF4070a (2.5 μ g/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with

citrate buffer pH 6, AP-staining.

cytokeratin 19 (aa285-298) Antibody (internal region) - References

Full-length cytokeratin-19 is released by human tumor cells: a potential role in metastatic progression of breast cancer. Alix-Panabières C, Vendrell JP, Slijper M, Pellé O, Barbotte E, Mercier G, Jacot W, Fabbro M, Pantel K. Breast cancer research : BCR 2009 11 (3): R39. PMID: 19549321