

**cytokeratin 19 (aa285-298), Biotinylated**  
Peptide-affinity purified goat antibody  
Catalog # AF4070b

**Specification**

---

**cytokeratin 19 (aa285-298), Biotinylated - Product Information**

|                   |  |
|-------------------|--|
| Application       | WB   |
| Primary Accession | <a href="#">P08727</a>                             |
| Other Accession   | <a href="#">NP_002267.2</a> , <a href="#">3880</a> |
| Reactivity        | Human  |
| Host              | Goat   |
| Clonality         | Polyclonal   |
| Concentration     | 0.5 mg/ml  |
| Isotype           | IgG  |
| Calculated MW     | 44106  |

**cytokeratin 19 (aa285-298), Biotinylated - Additional Information**

**Gene ID** 3880

**Other Names**

Keratin, type I cytoskeletal 19, Cytokeratin-19, CK-19, Keratin-19, K19, KRT19

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

cytokeratin 19 (aa285-298), Biotinylated is for research use only and not for use in diagnostic or therapeutic procedures.

**cytokeratin 19 (aa285-298), Biotinylated - Protein Information**

**Name** KRT19

**Function**

Involved in the organization of myofibers. Together with KRT8, helps to link the contractile apparatus to dystrophin at the costameres of striated muscle.

**Tissue Location**

Expressed in a defined zone of basal keratinocytes in the deep outer root sheath of hair follicles. Also observed in sweat gland and mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, ectocervical epithelium (at protein level). Expressed in epidermal basal cells, in nipple epidermis and a defined region of the


hair follicle. Also seen in a subset of vascular wall cells in both the veins and artery of human umbilical cord, and in umbilical cord vascular smooth muscle. Observed in muscle fibers accumulating in the costameres of myoplasm at the sarcolemma in structures that contain dystrophin and spectrin.

### **cytokeratin 19 (aa285-298), Biotinylated - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **cytokeratin 19 (aa285-298), Biotinylated - Images**



Biotinylated EB12359 (0.3µg/ml) staining of Human Colorectal cancer lysate (35µg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and usi

### **cytokeratin 19 (aa285-298), Biotinylated - References**

Full-length cytokeratin-19 is released by human tumor cells: a potential role in metastatic progression of breast cancer. Alix-Panabières C, Vendrell JP, Slijper M, Pellé O, Barbotte E, Mercier G, Jacot W, Fabbro M, Pantel K. Breast cancer research : BCR 2009 11 (3): R39. PMID: 19549321