

**PGK1 Antibody (internal region)**  
Peptide-affinity purified goat antibody  
Catalog # AF4083a

**Specification**

---

**PGK1 Antibody (internal region) - Product Information**

Application	WB
Primary Accession	<a href="#">P00558</a>
Other Accession	<a href="#">NP_000282.1</a> , <a href="#">5230</a>
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	44615

**PGK1 Antibody (internal region) - Additional Information**

**Gene ID** 5230

**Other Names**

Phosphoglycerate kinase 1, 2.7.2.3, Cell migration-inducing gene 10 protein, Primer recognition protein 2, PRP 2, PGK1, PGKA

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

PGK1 Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

**PGK1 Antibody (internal region) - Protein Information**

**Name** PGK1

**Synonyms** PGKA

**Function**

Catalyzes one of the two ATP producing reactions in the glycolytic pathway via the reversible conversion of 1,3- diphosphoglycerate to 3-phosphoglycerate (PubMed:<a href="http://www.uniprot.org/citations/30323285" target="\_blank">30323285</a>, PubMed:<a href="http://www.uniprot.org/citations/7391028" target="\_blank">7391028</a>). Both L- and D- forms of purine and pyrimidine nucleotides can be used as substrates, but the activity is much

lower on pyrimidines (PubMed:<a href="http://www.uniprot.org/citations/18463139" target="\_blank">18463139</a>). In addition to its role as a glycolytic enzyme, it seems that PGK1 acts as a polymerase alpha cofactor protein (primer recognition protein) (PubMed:<a href="http://www.uniprot.org/citations/2324090" target="\_blank">2324090</a>). Acts as a protein kinase when localized to the mitochondrion where it phosphorylates pyruvate dehydrogenase kinase PDK1 to inhibit pyruvate dehydrogenase complex activity and suppress the formation of acetyl- coenzyme A from pyruvate, and consequently inhibit oxidative phosphorylation and promote glycolysis (PubMed:<a href="http://www.uniprot.org/citations/26942675" target="\_blank">26942675</a>, PubMed:<a href="http://www.uniprot.org/citations/36849569" target="\_blank">36849569</a>). May play a role in sperm motility (PubMed:<a href="http://www.uniprot.org/citations/26677959" target="\_blank">26677959</a>).

#### Cellular Location

Cytoplasm, cytosol. Mitochondrion matrix. Note=Hypoxic conditions promote mitochondrial targeting (PubMed:26942675). Targeted to the mitochondrion following phosphorylation by MAPK1/ERK2, cis-trans isomerization by PIN1, and binding to mitochondrial circRNA mcPGK1 (PubMed:36849569).

#### Tissue Location


Mainly expressed in spermatogonia. Localized on the principle piece in the sperm (at protein level). Expression significantly decreased in the testis of elderly men

#### PGK1 Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### PGK1 Antibody (internal region) - Images



AF4083a (0.1 µg/ml) staining of HeLa lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

**PGK1 Antibody (internal region) - References**

Characterization of phosphoglycerate kinase-1 expression of stromal cells derived from tumor microenvironment in prostate cancer progression. Wang J, Ying G, Wang J, Jung Y, Lu J, Zhu J, Pienta KJ, Taichman RS. Cancer research 2010 Jan 70 (2): 471-80. PMID: 20068185