

CLIC1 / NCC27 (aa45-57) Antibody (internal region)
Peptide-affinity purified goat antibody
Catalog # AF4091a**Specification****CLIC1 / NCC27 (aa45-57) Antibody (internal region) - Product Information**

Application	WB
Primary Accession	O00299
Other Accession	NP_001279.2 , 1192 , 114584 (mouse) , 406864 (rat)
Reactivity	Human
Predicted	Mouse, Rat, Dog
Host	Goat
Clonality	Polyclonal
Concentration	0.5 mg/ml
Isotype	IgG
Calculated MW	26923

CLIC1 / NCC27 (aa45-57) Antibody (internal region) - Additional Information

Gene ID 1192

Other Names

Chloride intracellular channel protein 1, Chloride channel ABP, Nuclear chloride ion channel 27, NCC27, Regulatory nuclear chloride ion channel protein, hRNCC, CLIC1, G6, NCC27

Format

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CLIC1 / NCC27 (aa45-57) Antibody (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

CLIC1 / NCC27 (aa45-57) Antibody (internal region) - Protein Information

Name CLIC1 {ECO:0000303|PubMed:16339885, ECO:0000312|HGNC:HGNC:2062}

Function

In the soluble state, catalyzes glutaredoxin-like thiol disulfide exchange reactions with reduced glutathione as electron donor. Reduces selenite and dehydroascorbate and may act as an antioxidant during oxidative stress response (PubMed:25581026, PubMed:37759794). Can insert

into membranes and form voltage-dependent multi-ion conductive channels. Membrane insertion seems to be redox- regulated and may occur only under oxidizing conditions. Involved in regulation of the cell cycle.

Cellular Location

Nucleus. Nucleus membrane; Single-pass membrane protein. Cytoplasm. Cell membrane; Single-pass membrane protein. Endoplasmic reticulum {ECO:0000250|UniProtKB:Q6MG61}. Note=Mostly in the nucleus including in the nuclear membrane (PubMed:12681486, PubMed:9139710). Small amount in the cytoplasm and the plasma membrane (PubMed:9139710). Exists both as soluble cytoplasmic protein and as membrane protein with probably a single transmembrane domain (PubMed:11551966, PubMed:11940526, PubMed:12681486, PubMed:14613939, PubMed:9139710). Might not be present in the nucleus of cardiac cells (By similarity) {ECO:0000250|UniProtKB:Q6MG61, ECO:0000269|PubMed:11551966, ECO:0000269|PubMed:11940526, ECO:0000269|PubMed:12681486, ECO:0000269|PubMed:14613939, ECO:0000269|PubMed:9139710}

Tissue Location

Expression is prominent in heart, placenta, liver, kidney and pancreas.

CLIC1 / NCC27 (aa45-57) Antibody (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CLIC1 / NCC27 (aa45-57) Antibody (internal region) - Images



AF4091a (2 µg/ml) staining of U937 lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.

CLIC1 / NCC27 (aa45-57) Antibody (internal region) - References

Transmembrane extension and oligomerization of the CLIC1 chloride intracellular channel protein upon membrane interaction. Goodchild SC, Angstmann CN, Breit SN, Curmi PM, Brown LJ. *Biochemistry* 2011 Dec 50 (50): 10887-97. PMID: 22082111