

**Anti-14-3-3 sigma / Stratifin Antibody (Internal), Biotinylated**  
Catalog # AF4281a**Specification****Anti-14-3-3 sigma / Stratifin Antibody (Internal), Biotinylated - Product Information**

Application	WB
Primary Accession	<a href="#">P31947</a>
Other Accession	<a href="#">2810</a> , <a href="#">NP_006133.1</a> , <a href="#">55948</a> , <a href="#">313017</a>
Reactivity	Human, Mouse
Predicted	Human, Mouse, Rat, Pig, Dog
Calculated MW	27774

**Anti-14-3-3 sigma / Stratifin Antibody (Internal), Biotinylated - Additional Information****Gene ID** 2810**Other Names**

cancer; disease; epithelial marker

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Anti-14-3-3 sigma / Stratifin Antibody (Internal), Biotinylated is for research use only and not for use in diagnostic or therapeutic procedures.

**Anti-14-3-3 sigma / Stratifin Antibody (Internal), Biotinylated - Protein Information****Name** SFN**Synonyms** HME1 {ECO:0000303|PubMed:1390337}**Function**

Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathways (PubMed: [15731107](http://www.uniprot.org/citations/15731107), PubMed: [22634725](http://www.uniprot.org/citations/22634725), PubMed: [28202711](http://www.uniprot.org/citations/28202711), PubMed: [37797010](http://www.uniprot.org/citations/37797010)). Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif (PubMed: [15731107](http://www.uniprot.org/citations/15731107), PubMed: [22634725](http://www.uniprot.org/citations/22634725), PubMed: [28202711](http://www.uniprot.org/citations/28202711), PubMed: [37797010](http://www.uniprot.org/citations/37797010)). Binding generally results in the modulation of the activity of the binding partner (PubMed: [15731107](http://www.uniprot.org/citations/15731107), PubMed: [22634725](http://www.uniprot.org/citations/22634725), PubMed: [28202711](http://www.uniprot.org/citations/28202711), PubMed: [37797010](http://www.uniprot.org/citations/37797010)).

<http://www.uniprot.org/citations/22634725> target="\_blank">22634725</a>, PubMed:<a href="http://www.uniprot.org/citations/28202711" target="\_blank">28202711</a>, PubMed:<a href="http://www.uniprot.org/citations/37797010" target="\_blank">37797010</a>). Promotes cytosolic retention of GBP1 GTPase by binding to phosphorylated GBP1, thereby inhibiting the innate immune response (PubMed:<a href="http://www.uniprot.org/citations/37797010" target="\_blank">37797010</a>). Also acts as a TP53/p53-regulated inhibitor of G2/M progression (PubMed:<a href="http://www.uniprot.org/citations/9659898" target="\_blank">9659898</a>). When bound to KRT17, regulates protein synthesis and epithelial cell growth by stimulating Akt/mTOR pathway (By similarity). Acts to maintain desmosome cell junction adhesion in epithelial cells via interacting with and sequestering PKP3 to the cytoplasm, thereby restricting its translocation to existing desmosome structures and therefore maintaining desmosome protein homeostasis (PubMed:<a href="http://www.uniprot.org/citations/24124604" target="\_blank">24124604</a>). Also acts to facilitate PKP3 exchange at desmosome plaques, thereby maintaining keratinocyte intercellular adhesion (PubMed:<a href="http://www.uniprot.org/citations/29678907" target="\_blank">29678907</a>). May also regulate MDM2 autoubiquitination and degradation and thereby activate p53/TP53 (PubMed:<a href="http://www.uniprot.org/citations/18382127" target="\_blank">18382127</a>).

#### Cellular Location

Cytoplasm. Nucleus {ECO:0000250|UniProtKB:O70456} Secreted. Note=May be secreted by a non- classical secretory pathway.

#### Tissue Location

Present mainly in tissues enriched in stratified squamous keratinizing epithelium.


### Anti-14-3-3 sigma / Stratifin Antibody (Internal), Biotinylated - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-14-3-3 sigma / Stratifin Antibody (Internal), Biotinylated - Images





Biotinylated antibody (0.1  $\mu\text{g/ml}$ ) staining of Mouse Skin lysate (35  $\mu\text{g}$  protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.