

**Goat Anti-UBID4 Antibody (aa89-103) (internal region)**  
Catalog # AF4297a**Specification****Goat Anti-UBID4 Antibody (aa89-103) (internal region) - Product Information**

Application	WB
Primary Accession	<a href="#">Q92785</a>
Other Accession	<a href="#">NP_006259.1</a> , <a href="#">361711</a> , <a href="#">19708</a> , <a href="#">5977</a>
Reactivity	Human
Predicted	Human, Mouse, Rat, Pig, Dog
Host	Goat
Isotype	IgG
Calculated MW	44155

**Goat Anti-UBID4 Antibody (aa89-103) (internal region) - Additional Information**

Gene ID 5977

**Other Names**

Zinc finger protein ubi-d4, Apoptosis response zinc finger protein, BRG1-associated factor 45D, BAF45D, D4, zinc and double PHD fingers family 2, Protein requiem, DPF2, BAF45D, REQ, UBID4

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat Anti-UBID4 Antibody (aa89-103) (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat Anti-UBID4 Antibody (aa89-103) (internal region) - Protein Information**

Name DPF2

Synonyms BAF45D, REQ, UBID4

**Function**

Plays an active role in transcriptional regulation by binding modified histones H3 and H4 (PubMed: [27775714](http://www.uniprot.org/citations/27775714)), PubMed: [28533407](http://www.uniprot.org/citations/28533407)). Is a negative regulator of myeloid differentiation of hematopoietic progenitor cells (PubMed: [28533407](http://www.uniprot.org/citations/28533407)). Might also have a role in the development and maturation of lymphoid cells (By similarity). Involved in the regulation of non-canonical NF-kappa-B pathway (PubMed: [20460684](http://www.uniprot.org/citations/20460684)).

**Cellular Location**

Nucleus. Cytoplasm

**Tissue Location**


Ubiquitous.

**Goat Anti-UBID4 Antibody (aa89-103) (internal region) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Goat Anti-UBID4 Antibody (aa89-103) (internal region) - Images**



AF4297a (1  $\mu$ g/ml) staining of Jurkat (A) and K562 (B) lysate (35  $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.