

Goat Anti-UBID4 Antibody (aa89-103) (internal region)
Catalog # AF4297a**Specification****Goat Anti-UBID4 Antibody (aa89-103) (internal region) - Product Information**

Application	WB
Primary Accession	Q92785
Other Accession	NP_006259.1 , 361711 , 19708 , 5977
Reactivity	Human
Predicted	Human, Mouse, Rat, Pig, Dog
Host	Goat
Isotype	IgG
Calculated MW	44155

Goat Anti-UBID4 Antibody (aa89-103) (internal region) - Additional Information**Gene ID** 5977**Other Names**

Zinc finger protein ubi-d4, Apoptosis response zinc finger protein, BRG1-associated factor 45D, BAF45D, D4, zinc and double PHD fingers family 2, Protein requiem, DPF2, BAF45D, REQ, UBID4

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-UBID4 Antibody (aa89-103) (internal region) is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-UBID4 Antibody (aa89-103) (internal region) - Protein Information**Name** DPF2**Synonyms** BAF45D, REQ, UBID4**Function**

Plays an active role in transcriptional regulation by binding modified histones H3 and H4 (PubMed: [27775714](http://www.uniprot.org/citations/27775714)), PubMed: [28533407](http://www.uniprot.org/citations/28533407)). Is a negative regulator of myeloid differentiation of hematopoietic progenitor cells (PubMed: [28533407](http://www.uniprot.org/citations/28533407)). Might also have a role in the development and maturation of lymphoid cells (By similarity). Involved in the regulation of non-canonical NF-kappa-B pathway (PubMed: [20460684](http://www.uniprot.org/citations/20460684)).

Cellular Location

Nucleus. Cytoplasm

Tissue Location


Ubiquitous.

Goat Anti-UBID4 Antibody (aa89-103) (internal region) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-UBID4 Antibody (aa89-103) (internal region) - Images



AF4297a (1 μ g/ml) staining of Jurkat (A) and K562 (B) lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.