

Goat Anti-HOXA10 Antibody (internal region), Biotinylated
Catalog # AF4299a**Specification**

Goat Anti-HOXA10 Antibody (internal region), Biotinylated - Product Information

Application	WB
Primary Accession	P31260
Other Accession	NP_061824.3 , 3206
Reactivity	Human
Predicted	Human, Mouse, Pig
Host	Goat
Isotype	IgG
Calculated MW	42414

Goat Anti-HOXA10 Antibody (internal region), Biotinylated - Additional Information**Gene ID** 3206**Other Names**

Homeobox protein Hox-A10, Homeobox protein Hox-1.8, Homeobox protein Hox-1H, PL, HOXA10, HOX1H

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Goat Anti-HOXA10 Antibody (internal region), Biotinylated is for research use only and not for use in diagnostic or therapeutic procedures.

Goat Anti-HOXA10 Antibody (internal region), Biotinylated - Protein Information**Name** HOXA10**Synonyms** HOX1H**Function**

Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis. Binds to the DNA sequence 5'-AA[AT]TTTATTAC-3'.

Cellular Location


Nucleus.

Goat Anti-HOXA10 Antibody (internal region), Biotinylated - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Goat Anti-HOXA10 Antibody (internal region), Biotinylated - Images



Biotinylated AF4299a (0.2 µg/ml) staining of Human Skeletal Muscle lysate (35 µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.