

**Goat Anti-FBXO11 (aa754-765) Antibody**  
Peptide-affinity purified goat antibody  
Catalog # AF4327a

**Specification**

---

**Goat Anti-FBXO11 (aa754-765) Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q86XK2</a>
Other Accession	<a href="#">NP_079409.3</a> , <a href="#">NP_001177203.1</a> , <a href="#">XP_005264629.1</a> , <a href="#">XP_005264630.1</a> , <a href="#">XP_005264631.1</a> , <a href="#">XP_005264632.1</a>
Reactivity	Human
Predicted	Human, Mouse, Rat, Pig
Host	Goat
Clonality	Polyclonal
Concentration	100ug/200ul
Isotype	IgG
Calculated MW	103585

**Goat Anti-FBXO11 (aa754-765) Antibody - Additional Information**

**Gene ID** 80204

**Other Names**

F-box only protein 11, Protein arginine N-methyltransferase 9, Vitiligo-associated protein 1, VIT-1, FBXO11, FBX11, PRMT9, VIT1

**Format**

0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

**Immunogen**

Peptide with sequence C-KISSYTSYPMHD, from the internal region of the protein sequence according to [NP\\_079409.3](#); [NP\\_001177203.1](#); [XP\\_005264629.1](#); [XP\\_005264630.1](#); [XP\\_005264631.1](#); [XP\\_005264632.1](#).

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat Anti-FBXO11 (aa754-765) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat Anti-FBXO11 (aa754-765) Antibody - Protein Information**

**Name** FBXO11

## Synonyms FBX11, PRMT9, VIT1

### Function

Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins, such as DTL/CDT2, BCL6 and PRDM1/BLIMP1. The SCF(FBXO11) complex mediates ubiquitination and degradation of BCL6, thereby playing a role in the germinal center B-cells terminal differentiation toward memory B-cells and plasma cells. The SCF(FBXO11) complex also mediates ubiquitination and degradation of DTL, an important step for the regulation of TGF-beta signaling, cell migration and the timing of the cell-cycle progression and exit. Binds to and neddylates phosphorylated p53/TP53, inhibiting its transcriptional activity. Plays a role in the regulation of erythropoiesis but not myelopoiesis or megakaryopoiesis. Mechanistically, activates erythroid genes by mediating the degradation of BAHD1, a heterochromatin-associated protein that recruits corepressors to H3K27me3 marks (PubMed:<a href="http://www.uniprot.org/citations/33156908" target="\_blank">33156908</a>). Participates in macrophage cell death and inflammation in response to bacterial toxins by regulating the expression of complement 5a receptor 1/C5AR1 and IL-1beta (PubMed:<a href="http://www.uniprot.org/citations/33156908" target="\_blank">33156908</a>). Acts as a critical regulator to determine the level of MHC-II by mediating the recognition of degron at the P/S/T domain of CIITA leading to its ubiquitination and subsequent degradation via the proteasome (PubMed:<a href="http://www.uniprot.org/citations/37279268" target="\_blank">37279268</a>). Participates in the antiviral response by initiating the activation of TBK1-IRF3-IFN-I axis. Mediates the 'Lys-63'-linked ubiquitination of TRAF3 to strengthen the interaction between TRAF3 and TBK1 (PubMed:<a href="http://www.uniprot.org/citations/36897010" target="\_blank">36897010</a>).

### Cellular Location

Nucleus. Chromosome.

### Tissue Location

Isoform 5 is expressed in keratinocytes, fibroblasts and melanocytes.

## Goat Anti-FBXO11 (aa754-765) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Goat Anti-FBXO11 (aa754-765) Antibody - Images



