

**Goat anti-IFITM3 (N terminus) Antibody**  
Peptide-affinity purified goat antibody  
Catalog # AF4370a

**Specification**

---

**Goat anti-IFITM3 (N terminus) Antibody - Product Information**

Application	WB, IHC, Pep-ELISA
Primary Accession	<a href="#">Q01628</a>
Other Accession	<a href="#">NP_066362.2</a>
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Calculated MW	14632

**Goat anti-IFITM3 (N terminus) Antibody - Additional Information**

**Gene ID** 10410

**Other Names**

IFITM3; interferon induced transmembrane protein 3; 1-8U; DSPA2b; IP15; dispanin subfamily A member 2b; interferon-induced transmembrane protein 3; interferon-inducible protein 1-8U

**Format**

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

Goat anti-IFITM3 (N terminus) Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

**Goat anti-IFITM3 (N terminus) Antibody - Protein Information**

**Name** IFITM3 ([HGNC:5414](#))

**Function**

IFN-induced antiviral protein which disrupts intracellular cholesterol homeostasis. Inhibits the entry of viruses to the host cell cytoplasm by preventing viral fusion with cholesterol depleted endosomes. May inactivate new enveloped viruses which buds out of the infected cell, by letting them go out with a cholesterol depleted membrane. Active against multiple viruses, including influenza A virus, SARS coronaviruses (SARS-CoV and SARS-CoV-2), Marburg virus (MARV), Ebola virus (EBOV), Dengue virus (DENV), West Nile virus (WNV), human immunodeficiency virus type 1 (HIV-1), hepatitis C virus (HCV) and vesicular stomatitis virus (VSV) (PubMed:<a href="http://www.uniprot.org/citations/26354436" target="\_blank">26354436</a>, PubMed:<a href="http://www.uniprot.org/citations/26354436" target="\_blank">26354436</a>).

<http://www.uniprot.org/citations/33239446> target="\_blank">33239446</a>, PubMed:<a href="http://www.uniprot.org/citations/33270927" target="\_blank">33270927</a>). Can inhibit: influenza virus hemagglutinin protein- mediated viral entry, MARV and EBOV GP1,2-mediated viral entry, SARS- CoV and SARS-CoV-2 S protein-mediated viral entry and VSV G protein- mediated viral entry (PubMed:<a href="http://www.uniprot.org/citations/33270927" target="\_blank">33270927</a>). Plays a critical role in the structural stability and function of vacuolar ATPase (v-ATPase). Establishes physical contact with the v-ATPase of endosomes which is critical for proper clathrin localization and is also required for the function of the v-ATPase to lower the pH in phagocytic endosomes thus establishing an antiviral state. In hepatocytes, IFITM proteins act in a coordinated manner to restrict HCV infection by targeting the endocytosed HCV virion for lysosomal degradation (PubMed:<a href="http://www.uniprot.org/citations/26354436" target="\_blank">26354436</a>). IFITM2 and IFITM3 display anti-HCV activity that may complement the anti-HCV activity of IFITM1 by inhibiting the late stages of HCV entry, possibly in a coordinated manner by trapping the virion in the endosomal pathway and targeting it for degradation at the lysosome (PubMed:<a href="http://www.uniprot.org/citations/26354436" target="\_blank">26354436</a>). Exerts opposing activities on SARS-CoV-2, including amphipathicity-dependent restriction of virus at endosomes and amphipathicity-independent enhancement of infection at the plasma membrane (PubMed:<a href="http://www.uniprot.org/citations/33270927" target="\_blank">33270927</a>).

#### Cellular Location

Cell membrane; Single-pass type II membrane protein. Late endosome membrane; Single-pass type II membrane protein. Early endosome membrane; Single-pass type II membrane protein. Lysosome membrane; Single-pass type II membrane protein. Cytoplasm, perinuclear region. Note=Co-localizes with BRI3 isoform 1 at the perinuclear region.

#### Goat anti-IFITM3 (N terminus) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Goat anti-IFITM3 (N terminus) Antibody - Images