

**Goat anti-LIF Antibody**  
Peptide-affinity purified goat antibody  
Catalog # AF4527a

---

## Specification

### Goat anti-LIF Antibody - Product Information

Application	IHC, FC, Pep-ELISA
Primary Accession	<a href="#">P15018</a>
Other Accession	<a href="#">NP_002300.1</a>
Reactivity	Human
Host	Goat
Clonality	Polyclonal
Calculated MW	22008

### Goat anti-LIF Antibody - Additional Information

Gene ID 3976

#### Other Names

LIF; leukemia inhibitory factor (cholinergic differentiation factor); CDF; DIA; HILDA; D factor; MLPLI; OTTHUMP00000198698; cholinergic differentiation factor; differentiation inhibitory activity; differentiation stimulating factor; differentiation-induci

#### Format

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

Goat anti-LIF Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

### Goat anti-LIF Antibody - Protein Information

Name LIF

Synonyms HILDA

#### Function

LIF has the capacity to induce terminal differentiation in leukemic cells. Its activities include the induction of hematopoietic differentiation in normal and myeloid leukemia cells, the induction of neuronal cell differentiation, and the stimulation of acute-phase protein synthesis in hepatocytes.

#### Cellular Location

Secreted.

### **Goat anti-LIF Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Goat anti-LIF Antibody - Images**