

**Goat anti-Ku70 / G22P1 Antibody**  
Peptide-affinity purified goat antibody  
Catalog # AF4528a

### Specification

---

#### Goat anti-Ku70 / G22P1 Antibody - Product Information

Application	FC, Pep-ELISA
Primary Accession	<a href="#">P12956</a>
Other Accession	<a href="#">NP_001460.1</a> , <a href="#">NP_001275907.1</a>
Reactivity	Human, Mouse, Rat, Dog
Host	Goat
Clonality	Polyclonal
Calculated MW	69843

#### Goat anti-Ku70 / G22P1 Antibody - Additional Information

Gene ID 2547

#### Other Names

G22P1; thyroid autoantigen 70kDa (Ku antigen); HGNC:4055; CTC75; CTCBF; KU70; ML8; TLAA; ATP-dependent DNA helicase II, 70 kDa subunit; CTC box binding factor 75 kDa subunit; OTTHUMP00000028581; thyroid autoantigen 70kD (Ku antigen); thyroid-lupus autoant

#### Format

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin. Aliquot and store at -20°C. Minimize freezing and thawing.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

Goat anti-Ku70 / G22P1 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.

#### Goat anti-Ku70 / G22P1 Antibody - Protein Information

Name XRCC6

Synonyms G22P1

#### Function

Single-stranded DNA-dependent ATP-dependent helicase that plays a key role in DNA non-homologous end joining (NHEJ) by recruiting DNA-PK to DNA (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>, PubMed:<a

href="http://www.uniprot.org/citations/2466842" target="\_blank">2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">9742108</a>). Required for double-strand break repair and V(D)J recombination (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">9742108</a>). Also has a role in chromosome translocation (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">9742108</a>). Has a role in chromosome translocation (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">9742108</a>). The DNA helicase II complex binds preferentially to fork-like ends of double-stranded DNA in a cell cycle-dependent manner (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">9742108</a>). It works in the 3'-5' direction (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">9742108</a>). During NHEJ, the XRCC5-XRCC6 dimer performs the recognition step: it recognizes and binds to the broken ends of the DNA and protects them from further resection (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">9742108</a>). Binding to DNA may be mediated by XRCC6 (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">7957065</a>

target="\_blank">7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">9742108</a>). The XRCC5-XRCC6 dimer acts as a regulatory subunit of the DNA-dependent protein kinase complex DNA-PK by increasing the affinity of the catalytic subunit PRKDC to DNA by 100-fold (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">9742108</a>). The XRCC5-XRCC6 dimer is probably involved in stabilizing broken DNA ends and bringing them together (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">9742108</a>). The assembly of the DNA-PK complex to DNA ends is required for the NHEJ ligation step (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">9742108</a>). Probably also acts as a 5'-deoxyribose-5-phosphate lyase (5'-dRP lyase), by catalyzing the beta-elimination of the 5' deoxyribose-5-phosphate at an abasic site near double-strand breaks (PubMed:<a href="http://www.uniprot.org/citations/20383123" target="\_blank">20383123</a>). 5'-dRP lyase activity allows to 'clean' the termini of abasic sites, a class of nucleotide damage commonly associated with strand breaks, before such broken ends can be joined (PubMed:<a href="http://www.uniprot.org/citations/20383123" target="\_blank">20383123</a>). The XRCC5-XRCC6 dimer together with APEX1 acts as a negative regulator of transcription (PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>). In association with NAA15, the XRCC5-XRCC6 dimer binds to the osteocalcin promoter and activates osteocalcin expression (PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>). Plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP-RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway (PubMed:<a href="http://www.uniprot.org/citations/28712728" target="\_blank">28712728</a>).

#### Cellular Location

Nucleus. Chromosome

#### Goat anti-Ku70 / G22P1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Goat anti-Ku70 / G22P1 Antibody - Images**