

IgG (Immunoglobulin Gamma Heavy Chain) (B Cell Marker) Antibody - With BSA and Azide
Mouse Monoclonal Antibody [Clone SPM556]
Catalog # AH10499

Specification

IgG (Immunoglobulin Gamma Heavy Chain) (B Cell Marker) Antibody - With BSA and Azide - Product Information

Application	,14,3,4,
Primary Accession	P01857
Other Accession	3500 (IGHG1) , 3501 (IGHG2) , 3502 (IGHG3) , 3503 (IGHG4) , 510635 , P01859 , P01860 , P01861
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG2a, kappa
Calculated MW	75kDa KDa

IgG (Immunoglobulin Gamma Heavy Chain) (B Cell Marker) Antibody - With BSA and Azide - Additional Information

Other Names

Ig gamma-1 chain C region, IGHG1

Format

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

IgG (Immunoglobulin Gamma Heavy Chain) (B Cell Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

IgG (Immunoglobulin Gamma Heavy Chain) (B Cell Marker) Antibody - With BSA and Azide - Protein Information

Name IGHG1 {ECO:0000303|PubMed:11340299, ECO:0000303|Ref.13}

Function

Constant region of immunoglobulin heavy chains. Immunoglobulins, also known as antibodies, are membrane-bound or secreted glycoproteins produced by B lymphocytes. In the recognition phase of humoral immunity, the membrane-bound immunoglobulins serve as receptors which, upon binding of a specific antigen, trigger the clonal expansion and differentiation of B lymphocytes into immunoglobulins-secreting plasma cells. Secreted immunoglobulins mediate the effector phase of humoral immunity, which results in the elimination of bound antigens (PubMed:<a

<http://www.uniprot.org/citations/20176268> target="_blank">20176268, PubMed:22158414). The antigen binding site is formed by the variable domain of one heavy chain, together with that of its associated light chain. Thus, each immunoglobulin has two antigen binding sites with remarkable affinity for a particular antigen. The variable domains are assembled by a process called V-(D)-J rearrangement and can then be subjected to somatic hypermutations which, after exposure to antigen and selection, allow affinity maturation for a particular antigen (PubMed:17576170, PubMed:20176268). Mediates IgG effector functions on monocytes triggering ADCC of virus- infected cells.

Cellular Location


[Isoform 1]: Secreted

IgG (Immunoglobulin Gamma Heavy Chain) (B Cell Marker) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IgG (Immunoglobulin Gamma Heavy Chain) (B Cell Marker) Antibody - With BSA and Azide - Images



Formalin-fixed, paraffin-embedded human Tonsil stained with IgG Monoclonal Antibody (SPM556)

IgG (Immunoglobulin Gamma Heavy Chain) (B Cell Marker) Antibody - With BSA and Azide - Background

Recognizes a protein of 75kDa, identified as γ heavy chain of human immunoglobulins. It does not cross-react with α (IgA), μ (IgM), ϵ (IgE), or δ (IgD), heavy chains, T-cells, monocytes, granulocytes, or erythrocytes. This MAb is useful in the identification of leukemias, plasmacytomas, and certain non-Hodgkin's lymphomas. The most common feature of these malignancies is the restricted expression of a single heavy chain class. Demonstration of clonality in lymphoid infiltrates indicates

that the infiltrate is clonal and therefore malignant.