

Cyclin D1 (G1-Cyclin & Mantle Cell Marker) Antibody - With BSA and Azide
Mouse Monoclonal Antibody [Clone SPM587]
Catalog # AH10714

Specification

Cyclin D1 (G1-Cyclin & Mantle Cell Marker) Antibody - With BSA and Azide - Product Information

Application	,1,14,3,4,
Primary Accession	P24385
Other Accession	595 , 523852 , 667996
Reactivity	Human, Mouse, Rat, Monkey
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG2a, kappa
Calculated MW	36kDa KDa

Cyclin D1 (G1-Cyclin & Mantle Cell Marker) Antibody - With BSA and Azide - Additional Information

Gene ID 595

Other Names

G1/S-specific cyclin-D1, B-cell lymphoma 1 protein, BCL-1, BCL-1 oncogene, PRAD1 oncogene, CCND1, BCL1, PRAD1

Format

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

Cyclin D1 (G1-Cyclin & Mantle Cell Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

Cyclin D1 (G1-Cyclin & Mantle Cell Marker) Antibody - With BSA and Azide - Protein Information

Name CCND1 {ECO:0000303|PubMed:8204893, ECO:0000312|HGNC:HGNC:1582}

Function

Regulatory component of the cyclin D1-CDK4 (DC) complex that phosphorylates and inhibits members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle during G(1)/S transition (PubMed: [1827756](http://www.uniprot.org/citations/1827756) target="_blank">1827756, PubMed: [1833066](http://www.uniprot.org/citations/1833066) target="_blank">1833066, PubMed: [19412162](http://www.uniprot.org/citations/19412162) target="_blank">19412162, PubMed: [33854235](http://www.uniprot.org/citations/33854235) target="_blank">33854235)

target="_blank">33854235, PubMed:8114739, PubMed:8302605). Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase (PubMed:1827756, PubMed:1833066, PubMed:19412162, PubMed:8114739, PubMed:8302605). Hypophosphorylates RB1 in early G(1) phase (PubMed:1827756, PubMed:1833066, PubMed:19412162, PubMed:8114739, PubMed:8302605). Cyclin D-CDK4 complexes are major integrators of various mitogenic and antimitogenic signals (PubMed:1827756, PubMed:1833066, PubMed:19412162, PubMed:8302605). Also a substrate for SMAD3, phosphorylating SMAD3 in a cell-cycle-dependent manner and repressing its transcriptional activity (PubMed:15241418). Component of the ternary complex, cyclin D1/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex (PubMed:9106657). Exhibits transcriptional corepressor activity with INSM1 on the NEUROD1 and INS promoters in a cell cycle-independent manner (PubMed:16569215, PubMed:18417529).

Cellular Location

Nucleus. Cytoplasm Nucleus membrane. Note=Cyclin D-CDK4 complexes accumulate at the nuclear membrane and are then translocated to the nucleus through interaction with KIP/CIP family members


Cyclin D1 (G1-Cyclin & Mantle Cell Marker) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Cyclin D1 (G1-Cyclin & Mantle Cell Marker) Antibody - With BSA and Azide - Images





Formalin-fixed, paraffin-embedded human Mantle Cell Lymphoma stained with Cyclin D1 Ab (Clone SPM587).



Western Blot of Cyclin D1 in (1) C2C12, (2) HepG2, & (3) NIH3T3 Lysate with Cyclin D1 Ab (SPM587).

Cyclin D1 (G1-Cyclin & Mantle Cell Marker) Antibody - With BSA and Azide - Background

Recognizes a protein of 36kDa, identified as cyclin D1. Cyclin D1, one of the key cell cycle regulators, is a putative proto-oncogene overexpressed in a wide variety of human neoplasms. This antibody neutralizes the activity of cyclin D1 in vivo. About 60% of mantle cell lymphomas (MCL) contain a t(11; 14)(q13; q32) translocation resulting in over-expression of cyclin D1. This antibody is useful in identifying mantle cell lymphomas (cyclin D1 positive) from CLL/SLL and follicular lymphomas (cyclin D1 negative). Occasionally, hairy cell leukemia and plasma cell myeloma weakly express Cyclin D1.

Cyclin D1 (G1-Cyclin & Mantle Cell Marker) Antibody - With BSA and Azide - References

Lukas J, et. al. *Oncogene*, 1994, 9(3):707-18. | Gillett C, et. al. *Cancer Research*, 1994, 54(7):1812-7. | Bartkova J, et. al. *Journal of Pathology*, 1994, 172(3):237-45