

**gp100 / Melanosome / PMEL17 / SILV (Melanoma Marker) Antibody - With BSA and Azide  
Mouse Monoclonal Antibody [Clone SPM286 ]  
Catalog # AH10738**

**Specification**

**gp100 / Melanosome / PMEL17 / SILV (Melanoma Marker) Antibody - With BSA and Azide  
- Product Information**

Application	,14,3,4,
Primary Accession	<a href="#">P40967</a>
Other Accession	<a href="#">6490</a> , <a href="#">95972</a>
Reactivity	Human, Horse
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG2b, kappa
Calculated MW	90-100kDa KDa

**gp100 / Melanosome / PMEL17 / SILV (Melanoma Marker) Antibody - With BSA and Azide  
- Additional Information**

**Gene ID** 6490

**Other Names**

Melanocyte protein PMEL, ME20-M, ME20M, Melanocyte protein Pmel 17, Melanocytes lineage-specific antigen GP100, Melanoma-associated ME20 antigen, P1, P100, Premelanosome protein, Silver locus protein homolog, M-alpha, 95 kDa melanocyte-specific secreted glycoprotein, P26, Secreted melanoma-associated ME20 antigen, ME20-S, ME20S, M-beta, PMEL, D12S53E, PMEL17, SILV

**Format**

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

**Storage**

Store at 2 to 8°C. Antibody is stable for 24 months.

**Precautions**

gp100 / Melanosome / PMEL17 / SILV (Melanoma Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

**gp100 / Melanosome / PMEL17 / SILV (Melanoma Marker) Antibody - With BSA and Azide  
- Protein Information**

**Name** PMEL

**Synonyms** D12S53E, PMEL17, SILV

**Function**

Forms physiological amyloids that play a central role in melanosome morphogenesis and

pigmentation. The maturation of unpigmented premelanosomes from stage I to II is marked by assembly of processed amyloidogenic fragments into parallel fibrillar sheets, which elongate the vesicle into a striated ellipsoidal shape. In pigmented stage III and IV melanosomes, the amyloid matrix serves as a platform where eumelanin precursors accumulate at high local concentrations for pigment formation. May prevent pigmentation-associated toxicity by sequestering toxic reaction intermediates of eumelanin biosynthesis pathway.

#### **Cellular Location**

Endoplasmic reticulum membrane; Single-pass type I membrane protein. Golgi apparatus, cis-Golgi network membrane; Single-pass type I membrane protein. Endosome, multivesicular body. Melanosome Extracellular vesicle. Secreted. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065) Localizes predominantly to intraluminal vesicles (ILVs) within multivesicular bodies. Associates with ILVs found within the lumen of premelanosomes and melanosomes and particularly in compartments that serve as precursors to the striated stage II premelanosomes (PubMed:11694580, PubMed:12643545). Sorted to stage I melanosomes following its processing in the ER and cis-Golgi (PubMed:15096515) Transiently expressed at the cell surface before targeting to early melanosomes (PubMed:16760433, PubMed:30988362). Colocalizes with BACE2 in stage I and II melanosomes (PubMed:23754390). Colocalizes with CD63 and APOE at exosomes and in intraluminal vesicles within multivesicular endosomes (PubMed:21962903, PubMed:26387950)

#### **Tissue Location**


Normally expressed at low levels in quiescent adult melanocytes but overexpressed by proliferating neonatal melanocytes and during tumor growth. Overexpressed in melanomas. Some expression was found in dysplastic nevi.


### **gp100 / Melanosome / PMEL17 / SILV (Melanoma Marker) Antibody - With BSA and Azide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **gp100 / Melanosome / PMEL17 / SILV (Melanoma Marker) Antibody - With BSA and Azide - Images**





Formalin-fixed, paraffin-embedded human Melanoma stained with gp100 Monoclonal Antibody (SPM286).

**gp100 / Melanosome / PMEL17 / SILV (Melanoma Marker) Antibody - With BSA and Azide - Background**

By immunohistochemistry, it specifically recognizes a protein in melanocytes and melanomas. This MAb reacts with junctional and blue nevus cells and variably with fetal and neonatal melanocytes. Intradermal nevi, normal adult melanocytes, and non-melanocytic cells are negative. It does not stain tumor cells of epithelial, lymphoid, glial, or mesenchymal origin. This Mab labels formalin-fixed, paraffin-embedded melanomas and other tumors showing melanocytic differentiation.

**gp100 / Melanosome / PMEL17 / SILV (Melanoma Marker) Antibody - With BSA and Azide - References**

Vennegoor C. et al., Am.J. Pathol. 130: 179-192, 1988