

**Interferon alpha-2 (IFNA2) Antibody - With BSA and Azide**  
**Mouse Monoclonal Antibody [Clone N39 ]**  
**Catalog # AH11501**

**Specification**

---

**Interferon alpha-2 (IFNA2) Antibody - With BSA and Azide - Product Information**

|                   |   |
|-------------------|---|
| Application       | ,3,4,   |
| Primary Accession | <a href="#">P01563</a>                        |
| Other Accession   | <a href="#">3440</a> , <a href="#">211575</a> |
| Reactivity        | Human   |
| Host              | Mouse   |
| Clonality         | Monoclonal                                    |
| Isotype           | Mouse / IgG1, kappa                           |
| Calculated MW     | 16-27kDa KDa                                  |

**Interferon alpha-2 (IFNA2) Antibody - With BSA and Azide - Additional Information**

**Gene ID** 3440

**Other Names**

Interferon alpha-2, IFN-alpha-2, Interferon alpha-A, LeIF A, IFNA2, IFNA2A, IFNA2B, IFNA2C

**Storage**

Store at 2 to 8°C. Antibody is stable for 24 months.

**Precautions**

Interferon alpha-2 (IFNA2) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

**Interferon alpha-2 (IFNA2) Antibody - With BSA and Azide - Protein Information**

**Name** IFNA2

**Synonyms** IFNA2A, IFNA2B, IFNA2C

**Function**

Produced by macrophages, IFN-alpha have antiviral activities.

**Cellular Location**

Secreted.

**Interferon alpha-2 (IFNA2) Antibody - With BSA and Azide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Interferon alpha-2 (IFNA2) Antibody - With BSA and Azide - Images**

### **Interferon alpha-2 (IFNA2) Antibody - With BSA and Azide - Background**

Recognizes a protein of 16-27kDa, identified as human interferon-II) (IFN-(II). Its epitope maps between aa112-148 of IFN-II) (total aa172). This MAb is specific for IFN-(II) and does not cross-react with IFN-(I). The site recognized by this MAb is called **site I** and is responsible for the antiviral and anti-proliferative activities of IFN-(II). Epitopes of N27 and N39 MAbs are different and represent a good combination of antibodies to set up an ELISA assay for the quantitation of IFN-(II) after viral infections. The IFN- family consists of 24 or more genes or pseudo-genes. IFN-(II) is one of the two distinct families (I and II) of human IFN-. The -interferon are mainly produced by lymphocytes, monocytes, macrophages, and cell lines such as Namalwa and KG1 following induction by viruses, nucleic acids, and glucocorticoid hormones. They are involved in virus resistance on target cells, inhibition of cell proliferation, induction of cytokines and regulation of expression of MHC class I antigens.

### **Interferon alpha-2 (IFNA2) Antibody - With BSA and Azide - References**

Kontsek P et al. Mapping of two immunodominant structures on human interferon alpha 2c and their role in binding to cells. *Mol Immunol* 1991, 28:1289-1297 | Kontsek P et al. Peptide-mapping of three neutralizing epitopes into predicted biologically active sites of human interferon-alpha 2. *Immunol Lett* 1993, 35(3):281-284 | Pestka S et al. Interferons and their actions. *Annu Rev Biochem* 1987, 56:727-777 | Sen GC et al. The interferon system. A bird's eye view of its biochemistry. *J Biol Chem* 1992, 267(8):5017-5020 | Capon DJ et al. Two distinct families of human and bovine interferon-alpha genes are coordinately expressed and encode functional polypeptides. *Mol Cell Biol* 1985, 5(4):768-779 | Kurane I et al. Induction of interferon alpha from human lymphocytes by autologous, dengue virus-infected monocytes. *J Exp Med* 1987, 166(4):999-1010 | Lepe-Zuniga JL et al. Production of interferon-alpha induced by dsRNA in human peripheral blood mononuclear cell cultures: role of priming by dsRNA-induced interferons-gamma and -beta. *J Interferon Res* 1989, 9(4):445-456 | Aman MJ et al. Interferon-alpha stimulates production of interleukin-10 in activated CD4+ T cells and monocytes. *Blood* 1996, 87(11):4731-473