

**Insulin-like Growth Factor-1 (IGF-1) Antibody - With BSA and Azide**  
**Mouse Monoclonal Antibody [Clone IGF1/1020 ]**  
**Catalog # AH11509**

### Specification

---

**Insulin-like Growth Factor-1 (IGF-1) Antibody - With BSA and Azide - Product Information**

Application	,3,4,
Primary Accession	<a href="#">P05019</a>
Other Accession	<a href="#">3479</a> , <a href="#">160562</a>
Reactivity	Human, Mouse, Rat, Rabbit
Host	Mouse
Clonality	Monoclonal
Isotype	Mouse / IgG1, kappa
Calculated MW	~7.6kDa KDa

**Insulin-like Growth Factor-1 (IGF-1) Antibody - With BSA and Azide - Additional Information**

**Gene ID** 3479

**Other Names**

Insulin-like growth factor I, IGF-I, Mechano growth factor, MGF, Somatomedin-C, IGF1, IBP1

**Storage**

Store at 2 to 8°C. Antibody is stable for 24 months.

**Precautions**

Insulin-like Growth Factor-1 (IGF-1) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

**Insulin-like Growth Factor-1 (IGF-1) Antibody - With BSA and Azide - Protein Information**

**Name** IGF1 ([HGNC:5464](#))

**Function**

The insulin-like growth factors, isolated from plasma, are structurally and functionally related to insulin but have a much higher growth-promoting activity. May be a physiological regulator of [1-14C]- 2-deoxy-D-glucose (2DG) transport and glycogen synthesis in osteoblasts. Stimulates glucose transport in bone-derived osteoblastic (PyMS) cells and is effective at much lower concentrations than insulin, not only regarding glycogen and DNA synthesis but also with regard to enhancing glucose uptake. May play a role in synapse maturation (PubMed:<a href="http://www.uniprot.org/citations/21076856" target="\_blank">21076856</a>, PubMed:<a href="http://www.uniprot.org/citations/24132240" target="\_blank">24132240</a>). Ca(2+)-dependent exocytosis of IGF1 is required for sensory perception of smell in the olfactory bulb (By similarity). Acts as a ligand for IGF1R. Binds to the alpha subunit of IGF1R, leading to the activation of the intrinsic tyrosine kinase activity which autophosphorylates tyrosine residues in the beta subunit thus initiating a cascade of down-stream signaling events leading to activation of

the PI3K-AKT/PKB and the Ras-MAPK pathways. Binds to integrins ITGAV:ITGB3 and ITGA6:ITGB4. Its binding to integrins and subsequent ternary complex formation with integrins and IGFR1 are essential for IGF1 signaling. Induces the phosphorylation and activation of IGFR1, MAPK3/ERK1, MAPK1/ERK2 and AKT1 (PubMed:<a href="http://www.uniprot.org/citations/19578119" target="\_blank">19578119</a>, PubMed:<a href="http://www.uniprot.org/citations/22351760" target="\_blank">22351760</a>, PubMed:<a href="http://www.uniprot.org/citations/23243309" target="\_blank">23243309</a>, PubMed:<a href="http://www.uniprot.org/citations/23696648" target="\_blank">23696648</a>). As part of the MAPK/ERK signaling pathway, acts as a negative regulator of apoptosis in cardiomyocytes via promotion of STUB1/CHIP-mediated ubiquitination and degradation of ICER-type isoforms of CREM (By similarity).

#### **Cellular Location**

Secreted {ECO:0000250|UniProtKB:P05017}.

#### **Insulin-like Growth Factor-1 (IGF-1) Antibody - With BSA and Azide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **Insulin-like Growth Factor-1 (IGF-1) Antibody - With BSA and Azide - Images**

#### **Insulin-like Growth Factor-1 (IGF-1) Antibody - With BSA and Azide - Background**

This antibody is specific to Insulin-like Growth Factor (IGF-1) and shows minimal cross-reaction with IGF-11, Proinsulin, MSF, and Insulin. IGF-1 is a polypeptide growth factor with two isoforms that are produced by alternative splicing. Isoform 1 is also known as IGF-IB while isoform 2 is known as IGF-IA. IGF-1 stimulates the proliferation of a wide range of cell types including muscle, bone and cartilage tissue. It functions as an autocrine regulator of growth. Activation of IGF system has emerged as a key factor for tumor progression and resistance to apoptosis in many cancers like those of breast, thyroid and colon.

#### **Insulin-like Growth Factor-1 (IGF-1) Antibody - With BSA and Azide - References**

Rotwein p, et. al. (1986) J. Biol. Chem. 261: 4828-4832. | Sandberg-Nordqvist AC, et. al. (1993) Cancer Res. 53: 2475-2478. | Zheng WH, et. al. (2000) J. Neural Transm. Suppl. 2000: 261-272